



# Human exposure to indoor air pollutants in sleep microenvironments: A literature review



Brandon E. Boor<sup>a, b, \*</sup>, Michal P. Spilak<sup>c</sup>, Jelle Laverge<sup>d</sup>, Atila Novoselac<sup>e</sup>, Ying Xu<sup>e</sup>

<sup>a</sup> Lyles School of Civil Engineering, Purdue University, West Lafayette, IN, USA

<sup>b</sup> Ray W. Herrick Laboratories, Center for High Performance Buildings, Purdue University, West Lafayette, IN, USA

<sup>c</sup> Department of Civil, Architectural, and Environmental Engineering, Drexel University, Philadelphia, PA, USA

<sup>d</sup> Department of Architecture and Urban Planning, Ghent University, Ghent, Belgium

<sup>e</sup> Department of Civil, Architectural, and Environmental Engineering, Cockrell School of Engineering, The University of Texas at Austin, Austin, TX, USA

## ARTICLE INFO

### Article history:

Received 14 May 2017

Received in revised form

8 August 2017

Accepted 26 August 2017

Available online 30 August 2017

### Keywords:

Bedroom

Mattress dust

Allergens and microbes

Volatile and semi-volatile organic

compounds

Sleep quality

Human exposure

Particle resuspension

## ABSTRACT

We spend approximately one-third of our lives sleeping, yet little is known as to how human exposure to indoor air pollutants during sleep impacts human health and sleep quality. This paper provides a literature review of the current state-of-knowledge pertaining to human inhalation and dermal exposures while sleeping. An analysis of the duration of sleep exposure periods is provided, demonstrating that the sleep microenvironment is the predominant indoor space where humanity spends most of its time. Mattress dust is found to contain a diverse spectrum of biological particles and particle-bound chemical contaminants and their concentrations in dust can span many orders of magnitude among bed samples. These dust particles can become airborne through particle resuspension associated with body movements in bed. Mattress foam and covers, pillows, and bed frames can emit a variety of volatile and semi-volatile organic compounds, including phthalate plasticizers and organophosphate flame retardants, and emission rates can increase due to localized elevations in surface temperature and moisture near the bed due to close contact with the human body. This literature review demonstrates that human exposures to mattress-released pollutants can be amplified due to the source-proximity effect inherent to the sleep microenvironment, where the human body and breathing zone are in close and intimate contact with potential pollutant sources for prolonged periods. Given the findings of this review, human exposures to indoor air pollutants in the sleep microenvironment should receive more attention and future research is needed to fully understand how sleep exposures affect human health and sleep quality.

© 2017 Elsevier Ltd. All rights reserved.

## Contents

1. Introduction .....	529
2. Sleep exposure characteristics .....	529
2.1. Exposure pathways .....	529
2.2. Sleep exposure period .....	530
2.3. Exposure of infants and toddlers during sleep: the impact of body weight .....	531
3. Biological and chemical composition of mattress dust .....	531
3.1. House dust mite and animal allergens .....	531
3.2. Fungi .....	531
3.3. Bacteria .....	543
3.4. Plasticizers and flame retardants .....	543
3.5. Particle deposits on mattresses and bedding materials .....	544
4. Chemical emissions from mattresses, pillows, and bed/crib frames .....	544

\* Corresponding author. Hampton Hall of Civil Engineering, 550 Stadium Mall Drive, West Lafayette, IN, 47907, USA.

E-mail address: [bboor@purdue.edu](mailto:bboor@purdue.edu) (B.E. Boor).

4.1.	Volatile organic compounds (VOCs) .....	544
4.2.	Flame retardants .....	544
4.3.	Plasticizers .....	544
4.4.	Unreacted isocyanates .....	547
5.	The source-proximity effect of the sleep microenvironment .....	547
5.1.	Spatial proximity and concentration gradients near actively emitting sources .....	547
5.2.	Incomplete mixing of bedroom air .....	549
5.3.	Personal cloud effect and human-induced resuspension of mattress dust .....	549
5.4.	Body movements during sleep periods .....	549
5.5.	Human thermal plume .....	550
5.6.	Impact of body temperature and sweating on chemical emissions from mattresses .....	550
5.7.	Direct dermal contact with mattresses and bedding materials .....	550
5.8.	Personalized ventilation and air cleaning systems for beds .....	550
6.	Conclusions and future directions .....	551
	Acknowledgements .....	551
	References .....	551

## 1. Introduction

Many particulate and gaseous pollutants that are potentially health hazardous originate from indoor sources and concentrations are often much higher indoors than outdoors [133]. Since people spend about 80–90% of their time indoors [77,113,137], human exposures in the indoor environment are often substantially greater than exposures occurring outdoors. Across the world, adults typically sleep 8–9 h per day [46,77,114,159], virtually always indoors, which corresponds to one-third of their lifetime, making the sleep microenvironment particularly important in contributing to both their acute and chronic exposures to indoor pollutants. Mattresses, pillows, bedding materials, and bed frames are possible pollutant sources unique to sleep microenvironments. People are likely to be exposed to elevated concentrations of various chemical contaminants and biological particles since they are in close and intimate contact with these items. Early-life exposures to mattress-released pollutants are a particular concern, given the extended sleep periods of infants (12–14 h per day) and their low body weights [30,50,85,86].

The sleep microenvironment can be defined as the space encompassing a mattress, pillow, bedding materials, bed frame and the volume of air above these items that includes an individual's breathing zone (BZ) and buoyant thermal plume [15,16,88]. The seemingly innocuous sleep microenvironment can be home to a diversity of pollutants that have been shown to impact human health. Mite and animal allergens, human- and animal-associated bacteria, fungi, and semi-volatile organic compounds (SVOCs) can accumulate in settled dust on mattresses, pillows, and bed sheets (see references in Table 1). The materials used to manufacture mattresses and bedding products, such as polyurethane foam and vinyl mattress covers, are possible sources of a myriad of chemical contaminants, including volatile organic compounds (VOCs), plasticizers, flame retardants, and unreacted (free) isocyanates (NCO) (see references in Table 2).

Measuring indoor air pollutant concentrations at a reference location in the bulk air of a room may not be sufficient to characterize indoor exposures (e.g. Ref. [180]). This discrepancy, which is known as the source-proximity effect, is mainly due to the non-uniform distribution of pollutants that arises due to source location, the buoyant human thermal plume, occupant movements, and the overall airflow pattern and mixing conditions within a space [29,43,60,128,130,131]. The source-proximity effect is an important characteristic of the sleep microenvironment, where the human body and BZ are in close proximity to potential pollutant sources for

extended periods of time. Therefore, people may be exposed to elevated concentrations of pollutants via inhalation and dermal pathways as they sleep. Research on the underlying physical processes governing the source-proximity effect of the sleep microenvironment is limited.

Human exposure in sleep microenvironments is important, but has not been extensively researched in contrast to exposures in other types of indoor environments, such as classrooms, kitchens, and occupational workplaces. Therefore, it offers challenging research opportunities to advance our understanding of the pollutants commonly found in sleep microenvironments, the mechanisms by which pollutants are transported around the human body to an individual's BZ, pollutant concentrations and exposure levels that individuals experience while sleeping, and the total amount of pollutants that are inhaled or absorbed via dermal exposure. This knowledge is important to investigate subsequent health effects and to develop strategies to promote healthy bedrooms. As a key step in the advancement of this knowledge, we provide a literature review of the current understanding of human exposure in sleep microenvironments, including: exposure characteristics, a summary of the biological and chemical composition of mattress dust, an overview of chemical emissions from mattresses and bedding materials, the potential for elevated exposures due to the source-proximity effect, and the use of personalized ventilation and air cleaning systems for beds.

## 2. Sleep exposure characteristics

### 2.1. Exposure pathways

Inhalation exposure can be defined as the contact between an agent (pollutant) and a target (person) [179]. The contact between a pollutant and a person occurs at an exposure point, which we can define as a person's BZ. The inhalation exposure concentration can be defined as the mass of a pollutant (e.g. # of particles or  $\mu\text{g}$  of a VOC) in a person's BZ divided by the BZ volume. The volume of the BZ is variable, has been shown to be dependent on the transport dynamics of a specific pollutant e.g. Ref. [97], and is likely influenced by physical obstructions, such as a pillow for a person sleeping in the prone position [89]. However, it is generally defined as a hemisphere with a radius of 30–50 cm, with its center of origin at the mouth or the nose [72,132]. An expression for the time-integrated inhalation exposure,  $E_{ij}$ , of a specific pollutant,  $i$ , in a particular microenvironment,  $j$ , is:

$$E_{i,j} = \int_{t_0}^t C_{BZ,i,j}(t) dt \quad (1)$$

where  $C_{BZ,i,j}(t)$  is the BZ concentration at time  $t$ , although time-averaged BZ concentrations are commonly reported. Two key variables thus define inhalation exposures: the amount of pollutant that enters the BZ and the length of time over which continuous contact occurs between the pollutant and the BZ.

In addition to inhalation exposure assessment, it is necessary to evaluate transdermal uptake of chemical pollutants (e.g. SVOCs) in the sleep microenvironment, given the extended periods of contact between human skin and vinyl mattress covers, pillow protectors, and assorted bedding materials (bed sheets, blankets, duvet covers). Dermal exposures while sleeping can occur via both direct contact transfer with these items and air-to-skin uptake [169], the latter of which is linked to concentrations of gas-phase pollutants released from the mattress. Dermal exposure via direct contact transfer ( $E_{Dermal Contact}$ ) can be calculated using Equation (2):

$$E_{Dermal Contact} = C_s \cdot SA \cdot TE \cdot AF \quad (2)$$

where  $C_s$  is the surface concentration of a pollutant,  $SA$  is the contact area,  $TE$  is the efficiency of contact transfer, which represents the fraction of the surface pollutant that is transferred to the skin during a contact event, and  $AF$  is the fractional absorption factor. Although contact transfer has been well studied, numerical values of  $TE$  are difficult to measure accurately and thus remain poorly characterized. The value of  $TE$  can be influenced by the chemical compound, physical nature of the bedding surfaces, contact pressure and motion during sleep, sleeping posture, surface concentration of the SVOC, and temperature and humidity (e.g. Ref. [182]). Simulated experiments with full-size human thermal manikins could be used to determine empirical  $TE$ s for sleeping people.  $AF$  is typically an empirical quantity that is assumed to be a fixed value for a specific chemical regardless of exposure conditions. Although the  $AF$  approach has been widely adopted, Frasch et al. [183] discussed its potential limitations and concluded that loading, evaporation or sublimation of VOCs, and experimental duration, may have strong impacts on the fraction absorbed by skin.

Dermal exposure due to direct air-to-skin transport has received attention recently [9,167,169,170,175,181]. The transdermal uptake of a gas-phase pollutant ( $E_{Dermal Gas}$ ) can be estimated using Equation (3):

$$E_{Dermal Gas} = C_{gas} \cdot k_{p,g} \cdot BSA \quad (3)$$

where  $C_{gas}$  is the gas-phase concentration of a pollutant in proximity of the human body,  $k_{p,g}$  is the transdermal permeability coefficient, and  $BSA$  is the body surface area. Weschler and Nazaroff [169,181] demonstrated that the abundance of SVOCs on hand wipe samples can be predicted reasonably well from gas-phase concentrations using this expression. They developed equations to estimate  $k_{p,g}$  based on the knowledge of physical and chemical properties of indoor pollutants, such as molecular weight ( $MW$ ), octanol-water partition coefficient ( $K_{ow}$ ), and Henry's constant, and extended their analysis of transdermal penetration to approximately eighty VOC and SVOC compounds.

## 2.2. Sleep exposure period

The exposure period can be defined as the time of continuous contact between a pollutant and a person [179]. It is the magnitude of this period that makes the sleep microenvironment particularly

important in contributing to human inhalation and dermal exposures to various pollutants originating in mattresses, pillows, and bedding.

The U.S. Environmental Protection Agency (EPA) provides data on the duration of time spent (h/day) in a sleep or nap activity (EPA Exposure Factors Handbook (EFH) 2009). To investigate the dependence of sleep duration on age, we plotted the EFH's data, as shown in Fig. 1. During the first few months to years of life, infants (<1 year of age) and toddlers (1–3 years of age) spend a considerable amount of time sleeping, with an average of 13.3 h/day in the first year of life, 12.6 h/day in the second year, and 12.1 h/day in the third year, also reported in Refs. [37,70]. The data suggests that sleep microenvironments may play a critical role in characterizing exposures of very young children to indoor air pollutants. As expected, sleep duration continues to decline with age through much of adulthood, with an average of 8.2 h/day for the U.S. mean age group of 37 years [184]. An inflection is observed around 50 years of age, after which, sleep duration begins to steadily rise with age to an average of 9.2 h/day for people older than 81 years of age.

For adults, it is also constructive to evaluate how the time spent in the sleep microenvironment compares to other microenvironments in which people commonly spend their time. The National Human Activity Pattern Survey (NHAPS) Study conducted by Ref. [77] surveyed over 9000 adults (>18 years of age) in the U.S. on their activity patterns, finding that Americans spent nearly 87% of their time indoors. The NHAPS data can be used along with the EFH data set to estimate the average percentage of time an adult in the U.S. spends in sleep microenvironments. We assumed that the time spent in the sleep or nap activity (mean of 8.2 h for adult of mean age in the U.S.), as defined by the U.S. EPA, is spent entirely in a person's bedroom, on their mattress or similar bedding, and in their personal residence. As shown in Fig. 2, the modified activity pattern data shows us that adults spend about 34% of their day in the sleep microenvironment, which equates to about 50% of the time they spend in a residence, and 39% of the time they spend indoors. Assuming the "in a residence, other activity" location category can be partitioned among several different microenvironments, such as living areas, kitchen, office, and bathroom, the sleep microenvironment becomes the predominant indoor space where American adults spend most of their time. Furthermore, the percentage of time spent in sleep microenvironments is likely an underestimate, given that the U.S. EPA EFH data is limited to time spent sleeping or napping, and does not include other activities common to this indoor space, such as reading a book, watching television in bed, or using a smartphone/tablet.

Time-activity pattern studies conducted outside of the U.S. have reported similar trends in time spent sleeping. Among Canadian adults, 36% of the day is spent sleeping or napping, predominately in one's residence [92]. In Europe, adult sleeping time varies by country, but is generally between 8 and 9 h/day: Belgium: 8.4 h/day; Bulgaria: 9.1 h/day; Estonia: 8.4 h/day; Spain: 8.6 h/day; France: 8.9 h/day; Italy: 8.3 h/day; Latvia: 8.7 h/day; Lithuania: 8.5 h/day; Poland: 8.5 h/day; Slovenia: 8.4 h/day; Finland: 8.5 h/day; United Kingdom: 8.4 h/day; and Norway: 8 h/day [46]. Adults in Australia and New Zealand spend 8.5 h/day sleeping; in China adults sleep 9 h/day; while in Japan and the Republic of Korea, adults sleep less than 8 h/day [114]. Furthermore, there is additional variability in the time spent indoors and in the home among different countries due to differences in climate and culture, e.g. Refs. [113,137,177]. Notably, Koreans spend 59% and 67% of their time indoors on weekdays and weekends, respectively, much less than North Americans and Europeans [177].

### 2.3. Exposure of infants and toddlers during sleep: the impact of body weight

Because of their low body weight, infants and toddlers inhale considerably more air per kg of body weight as they sleep compared to adolescents and adults. The volume of air inhaled/kg-day can be estimated with the U.S. EPA EFH data set by taking the product of the mean normalized volumetric breathing rate in the sleep or nap activity (L/h-kg) and the mean duration of time spent in the sleep or nap activity (h/day). The normalized inhaled air volumes,  $V_{Sleep}^*$ , are categorized by age group and gender and presented in Fig. 3. It is apparent that infants and toddlers inhale nearly an order of magnitude more air per body mass than adults. During the first and second years of life, they inhale approximately 300 L/kg-day, which decreases to about 244 L/kg-day during the third year of life. Between 21 and < 31 years of age, adults breathe about 30 L/kg-day.

The analysis above helps to understand how the dose of an inhaled pollutant is considerably different between very young children and adults. The daily inhalation intake dose,  $D_{i,j}$ , (mass of pollutant inhaled per body weight per day) of pollutant  $i$  in microenvironment  $j$  can be calculated as:

$$D_{i,j} = \frac{\int_{t_0}^t Q_{B,j}(t) \times C_{BZ,i,j}(t) dt}{BW} \quad (4)$$

where  $Q_{B,j}(t)$  is the volumetric breathing rate (L/h) and  $BW$  is body weight (kg). This expression can be simplified for the sleep microenvironment and by assuming time-averaged BZ concentrations and breathing rates. Additionally, the normalized inhaled air volumes during a sleep period,  $V_{Sleep}^*$ , can be substituted into the equation to obtain:

$$\overline{D_{i,Sleep}} = \frac{\overline{Q_{B,Sleep}} \times t_{Sleep} \times \overline{C_{BZ,i,Sleep}}}{BW} = V_{Sleep}^* \times \overline{C_{BZ,i,Sleep}} \quad (5)$$

Thus, to estimate the daily sleeping dose, one can simply take the product of  $V_{Sleep}^*$ , as calculated from the U.S. EPA EFH data set, and the average BZ concentration of a pollutant originating in the sleep microenvironment. Therefore, if a very young child (an infant or toddler) and an adult are exposed to the same BZ concentration of a pollutant released from a mattress, the child-normalized dose will be an order of magnitude greater than that of the adult. Similarly, to compare an infant's dose of dermally absorbed pollutants to that of an adult, we can use the skin surface area to body mass ratio ( $BSA/BW$ ), as presented in Fig. 4. Infants have  $BSA/BW$  three times greater than adults, emphasizing the importance of early-life dermal exposures, particularly during sleep periods.

### 3. Biological and chemical composition of mattress dust

Mattresses, pillows, and bedding materials serve as an accumulation zone for a diverse spectrum of biological particles and particle-bound chemical contaminants. Table 1 provides an overview of the biological and chemical composition of mattress dust in selected field studies ( $n = 52$ ). These studies are often aimed at identifying associations between the microbial and allergenic content of mattress dust with the development of atopic and allergic diseases in infants and children, e.g. Refs. [21,27,80,125,146,154,160]. Mattress dust is commonly sampled as a surrogate for human exposure in large cohort studies. However, air sampling, particularly near the BZ, may provide more valuable insight in characterizing sleep inhalation exposures, e.g. Refs. [57,102,122,134], as dust levels are not necessarily correlated

with concentrations in the air. However, field data on BZ concentrations during sleep is limited.

#### 3.1. House dust mite and animal allergens

Biological matter in mattress dust is comprised of a wide range of organisms and their associated allergens. Among house dust mite (HDM) allergens, Der p 1, Der f 1, Der p2, and Blo t 5, have been identified in many regions of the world across a range of climate zones. The high moisture levels and plentiful supply of desquamated skin particles, or squames, makes the sleep microenvironment an ideal environment for HDMs to flourish. For example, in Brazil, da Silva et al. [35] detected 400–1000 HDM bodies (eggs, larvae, nymphs, and adults) per g of mattress dust. Additionally, cat (Fel d 1), dog (Can f 1), cockroach (Bla g 2), and mouse/rat urinary allergens have also been detected, along with allergens originating outdoors (Ph1 p 5, timothy pollen). Concentrations of these allergens can range over several orders of magnitude, from <1 to > 10<sup>3</sup> µg/g, as shown in Table 1.

The type of mattress and bedding materials has been shown to influence levels of HDM allergens. Van den Bemt et al. [161] reported higher levels on inner spring mattresses, compared to polyester and latex, and waterbeds. In Thailand, [163] found varying levels Der p 1 and Der f 1 in different mattresses, with kapok showing the highest concentrations, followed (in decreasing order) by synthetic, coconut, and polyurethane sponge. Der p 1 levels also tend to increase with mattress age [107,161]. In addition, the frequent use of wool, synthetic, and sheepskin bedding materials was found to be associated with higher levels of Der p 1 allergens [51,107]; and pillow dust can also be a source of Der p 1, Der f 1, and Blo t 5 allergens [172]. Finally, differences in HDM allergen levels have also been observed between the mattresses of mothers and their infants [26], between rural and urban residences [44], before and after home renovation [200], as well as variations among the seasons [150].

Additional studies have reported elevated levels of airborne HDM allergens during sleep periods, suggesting that deposited HDM particles can be released into the air via human-induced resuspension, e.g. Refs. [16,123,141]. Gore et al. [57] measured airborne Der p 1 allergen exposure via nasal air sampling and estimated inhalation exposures of 6 ng/night and 2 µg/year during sleep periods. Sakaguchi et al. [134] reported airborne Der p 1/Der f 1 and Der 2 concentrations of 223 and 87.1 pg/m<sup>3</sup>, respectively, in the air directly above a mattress. These levels were found to be a factor of ten greater than those measured in the living room air of the same residence, which may suggest significant resuspension of settled HDM particles from mattresses. However, a recent study has suggested that beds may not be the primary site of HDM exposure [156].

#### 3.2. Fungi

A wide variety of fungal genera and species are commonly identified in mattress dust, including *Penicillium* spp., *Cladosporium* spp., *Aspergillus* spp., *Aspergillus fumigatus* spp., *Alternaria* spp., *Eurotium* spp., *Epicoccum*, and yeasts, among many others (Table 1). Along with dust, many fungal species have been found within mattress and pillow foam, with early studies by Refs. [31,33] recognizing the impact of fungal contamination of pillows on human health. Concentrations of fungal species are typically reported as CFU (colony forming units) per g of mattress dust or number of cells per g of mattress dust, with concentrations in the range of 10<sup>3</sup>–10<sup>4</sup> CFU/g and < 10<sup>4</sup> to > 10<sup>7</sup> cells/g, respectively. Studies also report levels of (1–3)-β-Glucan, a component of the cell wall of many fungi (also found in some bacteria) and a common microbial

**Table 1**  
Biological material and chemical contaminants detected in mattress dust in selected field studies ( $n = 52$ ).

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
1. Garrett et al. (1998) [51]	Evaluated the impact of bedding characteristics on HDM concentrations	Latrobe Valley, Victoria, Australia	Der p 1	<b>Wool bedding:</b> Always: 30.6 (24.8–37.8) Sometimes: 23.6 (20.0–27.7) Never: 11.2 (8.6–14.6) <b>Type of mattress:</b> Inner-spring: 25.0 (21.9–28.6) Foam: 13.0 (8.9–19.1)	µg/g mattress dust (Mean (95% CI))	Not reported
2. Douwes et al. (1998) [38]	Relationship between house characteristics and levels of endotoxin and (1–3)-β-D-Glucan	Erfurt and Hamburg, Germany	Endotoxin (1–3)-β-D-Glucan	Hamburg: 300 Erfurt: 4200 Hamburg: 317 Erfurt: 862	EU/m <sup>2</sup> mattress area sample (Mean) µg/m <sup>2</sup> mattress area sampled (Mean)	0.528 (Hamburg) 1.075 (Erfurt) (Mean)
3. Fahlbusch et al. (2000) [47]	Quantified indoor grass pollen allergens in house dust	Hamburg, Erfurt, Zerbst, Bitterfeld & Hettstedt, Germany	Phl p 5	<0.03 (<0.03–0.120)	µg/g mattress dust (Median (25 <sup>th</sup> -75 <sup>th</sup> percentiles))	Not reported
4. Stubner et al. (2000) [149]	Examined the impact of home remediation interventions on lowering allergen levels in low-income households	Birmingham, Alabama, USA	Various allergens (e.g. Der p 1, Der f 1, Fel d 1)	<b>Home 1, Mattress:</b> 15 (before intervention), <2 (after intervention) <b>Home 2, Mattress:</b> 20 (before intervention), 4 (after intervention) <b>Home 3, Mattress:</b> 235 (before intervention), 16 (after intervention) <b>Home 4, Mattress:</b> 840 (before intervention), – (after intervention) <b>Home 5, Mattress 1:</b> 630 (before intervention), 240 (after intervention) <b>Home 5, Mattress 2:</b> 565 (before intervention), – (after intervention) <b>Home 6, Mattress:</b> 5 (before intervention), – (after intervention)	µg/mg mattress dust	Not reported
5. Hirsch et al. (2000) [200]	Measured HDM and mold spore concentrations before and after installation of windows and central heating systems	Dresden, Germany	Der f 1 <i>Cladosporium</i> spp. <i>Penicillium</i> spp. <i>Alternaria</i> spp. <i>Aspergillus</i> spp. <i>Aspergillus fumigatus</i> spp.	<b>Before renovation:</b> 1.56 (0.89–2.74) <b>After renovation:</b> 2.40 (1.33–4.36) <b>Before renovation:</b> 130 (50–370) <b>After renovation:</b> 60 (20–160) <b>Before renovation:</b> 1500 (740–3020) <b>After renovation:</b> 1250 (590–2660) <b>Before renovation:</b> 10 (0–20) <b>After renovation:</b> 10 (10–30) <b>Before renovation:</b> 590 (270–1320) <b>After renovation:</b> 820 (400–1710) <b>Before renovation:</b> 10 (0–10) <b>After renovation:</b> 10 (10–20)	µg/g mattress dust (Mean (95% CI)) CFU/g mattress dust (Mean (95% CI))	Not reported

6. Douwes et al. (2000) [39]	Investigated the relationship between endotoxin, $\beta$ -Glucans, and HDM allergens in mattress dust with peak expiratory flow in children	Amsterdam, Netherlands	Endotoxin	<b>Non-symptomatics:</b> 1820 $\pm$ 3.6 (level), 4772 $\pm$ 2.8 (load) <b>Symptomatics:</b> 2082 $\pm$ 4.3 (level), 5696 $\pm$ 3.6 (load) <b>Asthmatics:</b> 1202 $\pm$ 3.3 (level), 3983 $\pm$ 1.8 (load) <b>Cough:</b> 3402 $\pm$ 4.7 (level), 7670 $\pm$ 4.3 (load)	Level: EU/m <sup>2</sup> mattress area sampled Load: EU/g mattress dust (Mean $\pm$ SD)	Not reported
			(1–3)- $\beta$ -D-Glucan	<b>Non-symptomatics:</b> 276 $\pm$ 2.4 (level), 718 $\pm$ 1.8 (load) <b>Symptomatics:</b> 293 $\pm$ 2.3 (level), 792 $\pm$ 2.1 (load) <b>Asthmatics:</b> 283 $\pm$ 1.9 (level), 903 $\pm$ 1.8 (load) <b>Cough:</b> 303 $\pm$ 2.6 (level), 683 $\pm$ 2.3 (load)	Level: $\mu$ g/m <sup>2</sup> mattress area sampled Load: $\mu$ g/g mattress dust (Mean $\pm$ SD)	
			Der p 1	<b>Non-symptomatics:</b> 537 $\pm$ 6.1 (level), 1440 $\pm$ 6.1 (load) <b>Symptomatics:</b> 432 $\pm$ 7.5 (level), 1147 $\pm$ 6.4 (load) <b>Asthmatics:</b> 318 $\pm$ 9.0 (level), 960 $\pm$ 7.8 (load) <b>Cough:</b> 646 $\pm$ 5.9 (level), 1481 $\pm$ 5.9 (load)	Level: ng/m <sup>2</sup> mattress area sampled Load: ng/g mattress dust (Mean $\pm$ SD)	
7. Su et al. (2001) [150]	Measured seasonal concentrations of HDM allergens and endotoxin	Southern Taiwan	Der p 1	<b>Spring:</b> 3.78, <b>Summer:</b> 5.61, <b>Fall:</b> 9.11, <b>Winter:</b> 10.15	$\mu$ g/g mattress dust (Mean)	Not reported
			Der p 2	<b>Spring:</b> 1.41, <b>Summer:</b> 1.49, <b>Fall:</b> 1.96, <b>Winter:</b> 3.32		
8. Pitten et al. (2001) [119]	Investigated the impact of mattress encasings on fungal growth	Germany	Endotoxin	<b>Spring:</b> 45.18, <b>Summer:</b> 17.96, <b>Fall:</b> 50.25, <b>Winter:</b> 18.84	ng/mg mattress dust (Mean)	Not reported
			<b>Fungal genera and species identified in dust on mattresses with cotton encasings:</b> <i>Alternaria</i> spp., <i>Aspergillus</i> spp., <i>Candida</i> spp., <i>Cladosporium cladosporioides</i> , <i>Mycelia sterilia</i> , <i>Penicillium</i> spp., <i>Rhodotorula</i> spp., <i>Scopulariopsis brevicaulis</i> , <i>Scopulariopsis</i> spp., <i>Sirodesmium-like fungus</i> , <i>Trichoderma</i> spp., <i>Ulocladium</i> spp. <b>Fungal genera and species identified in dust on mattresses with synthetic encasings:</b> <i>Aspergillus</i> spp., <i>Cladosporium cladosporioides</i> , <i>Mycelia sterilia</i> , <i>Penicillium</i> spp., <i>Scopulariopsis brevicaulis</i> , <i>Scopulariopsis</i> spp.			
9. Oppermann et al. (2001) [117]	Measured levels of HDM allergens and fungi in children's mattresses	East & West Germany	Der f 1	1.36	$\mu$ g/g mattress dust (Mean)	Not reported
			Der p 1	1.14		
10. Mahrshahi et al. (2002) [107]	Measured HDM allergens in residential homes and investigated the impact of mattress and bedding characteristics on concentrations	Sydney, Australia	Common fungal species identified:	<b>Total fungi:</b> 26,500 (1400–300,000)	CFU/g mattress dust (Mean (Range))	Not reported
			Der p 1	<b>Type of mattress:</b> Foam: 21.39 (13.77–33.25) Inner spring: 14.22 (12.58–16.07) <b>Mattress Age:</b> $\leq$ 2 years: 10.31 (8.03–13.24) >2 years: 15.47 (13.47–17.67) <b>Wool blankets:</b> Yes: 27.18 (21.06–35.08) No: 12.88 (11.38–14.57) <b>Synthetic blankets:</b> Yes: 22.18 (17.51–28.11) No: 13.21 (11.64–14.99) <b>Sheepskin:</b> Yes: 22.95 (14.37–36.66) No: 13.84 (12.31–15.56)	$\mu$ g/g mattress dust (Mean (95% CI))	

Table 1 (continued)

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
11. Sidenius et al. (2002) [138]	Distribution of HDMs and their allergens on different surfaces in houses occupied by patients with newly diagnosed allergy to HDM allergen and previously reported high Der 1 concentrations in mattresses	Copenhagen, Denmark	Der 1	86 (30–288) µg/g; 88 (12–1910) mites/g. Note: 100% of samples from mattresses exceeded 10 µg/g; 37.5% of samples from pillows and duvets exceeded 10 µg/g	µg/g mattress dust; mites/g mattress dust (Median)	Not reported
12. Jovanovic et al. (2004) [73]	Measured indoor fungi levels in homes of children with and without a history of allergies	Southwest Germany	Common fungal species identified: <i>Aspergillus</i> spp., <i>Penicillium</i> spp., <i>Alternaria</i> spp., <i>Cladosporium</i> spp.	<b>Total viable mold spores:</b> 16,250/38,238 (0–2,500,000)	CFU/g mattress dust (Median/Mean (Range))	Children with allergic history: 0.393 Controls: 0.403 (Median)
13. Gehring et al. (2004) [52]	Measured endotoxin concentrations in the mattress dust of mothers and their infants	Munich & Leipzig, Germany	Endotoxin	<b>Mothers' mattress:</b> Munich: 2000 (level), 3000 (load) Leipzig: 2200 (level), 3100 (load) <b>Infants' mattress:</b> Munich: 900 (level), 4900 (load) Leipzig: 1000 (level), 6600 (load)	Level: EU/m <sup>2</sup> mattress area sampled Load: EU/g mattress dust (~Mean)	Infants' mattress: ~0.2 Mothers' mattress: ~0.7
14. El Sharif et al. (2004) [44]	Measured concentrations of HDM allergens, pet allergens, and endotoxin in mattress dust	Ramallah, Palestine	Der p 1	<b>City residences:</b> 958 (load), 4290 (level) <b>Village residences:</b> 1941 (load), 3776 (level) <b>Camp residences:</b> 3100 (load), 5656 (level)	Load: ng/g mattress dust Level: ng/m <sup>2</sup> mattress area sampled (Mean)	City residences: 0.29 ± 0.20 Village residences: 0.59 ± 0.31 Camp residences: 0.62 ± 0.36
			Der f 1	<b>City residences:</b> 48 (load), 206 (level) <b>Village residences:</b> 33 (load), 64 (level) <b>Camp residences:</b> 24 (load), 42 (level)		
			Can f 1	<b>City residences:</b> 100 (load), 451 (level) <b>Village residences:</b> 99 (load), 192 (level) <b>Camp residences:</b> 164 (load), 298 (level)		
			Fel d 1	<b>City residences:</b> 91 (load), 408 (level) <b>Village residences:</b> 210 (load), 408 (level) <b>Camp residences:</b> 151 (load), 275 (level)		
			Endotoxin	<b>City residences:</b> - (load), 18,738 (level) <b>Village residences:</b> 19,214 (load), 25,700 (level) <b>Camp residences:</b> 23,972 (load), 43,696 (level)	Load: EU/g mattress dust Level: EU/m <sup>2</sup> mattress area sampled (Mean)	
15. Instanes et al. (2005) [71]	Measured concentrations of allergens and endotoxin in mattress dust in day-care centers	Oslo, Norway	Fel d 1 Can f 1	1.5 (load), 0.17 (level) 10.0 (load), 1.7 (level)	Load: µg/g mattress dust Level: µg/m <sup>2</sup> mattress area sampled (Median)	Not reported

			Endotoxin	5.0 (load), 0.9 (level)	Load: ng/g mattress dust Level: ng/m <sup>2</sup> mattress area sampled (Median)	
16. da Silva et al. (2005) [35]	Investigated mite fauna in mattress dust	Londrina, Brazil	HDM bodies: total mites  HDM bodies: Pyroglyphidae mites	<b>Crib:</b> 404.5 ± 183.3 <b>Bed:</b> 1075.8 ± 198.8 <b>Crib:</b> 289.9 ± 136.7 <b>Bed:</b> 875.0 ± 183.6	Mite bodies/g mattress dust (Mean ± 95% CI) Note: bodies represent all life stages: eggs, larvae, nymphs, and adults.	Not reported
17. Hicks et al. (2005) [63]	Analyzed fungal flora and concentrations in settled dust	Northern California, USA	Der p 1  Culturable fungi  <i>Cladosporium</i> <i>Penicillium</i> spp. Yeasts <i>Aureobasidium</i> <i>Aspergillus niger</i> Nonsporulating fungi <i>Alternaria</i> <i>Epicoccum</i>	<b>Crib:</b> 5.8 ± 2.7 <b>Bed:</b> 17.5 ± 3.7 62,300 (37,300–104,000)  22,000 (ND–180,000) 4000 (ND–120,000) 8800 (NS–200,000) 4300 (ND–68,000) 400 (ND–56,000) 800 (ND–1400) 1800 (ND–12,000) 800 (ND–12,000)	µg/g mattress dust (Mean ± 95% CI, estimate) CFU/g bedspread and furniture dust (Mean (95% CI)) CFU/g bedspread and furniture dust (Median (Range))	Not reported
18. van den Bemt et al. (2006) [161]	Evaluated the influence of mattress characteristics on HDM concentrations	Netherlands	Der p 1	<b>Type of mattress:</b> Latex: 1.191 (0.744–1.908) Waterbed: 0.433 (0.073–2.559) Polyester: 1.199 (0.592–2.431) Inner spring: 1.418 (0.594–3.387) <b>Upper layer material:</b> Cotton: 0.801 (0.343–1.873) Synthetic: 2.606 (1.527–4.448) Cotton/synthetic: 1.382 (0.885–2.158) Other: 0.447 (0.111–1.792) <b>Mattress age:</b> ≤5 years: 0.986 (0.517–1.883) >5 years: 1.190 (0.731–1.939) <b>Relative humidity in bedroom:</b> ≤50%: 0.796 (0.444–1.426) >50%: 1.975 (1.312–2.974)	log µg/g mattress dust (Mean (95% CI))	Not reported
19. Woodcock et al. (2006) [171]	Identified fungal flora in used (18 months to >20 years) synthetic and feather pillows	United Kingdom	<i>Penicillium</i> spp. <i>Aspergillus vitus</i> <i>Rhodotorula mucilaginosa</i> <i>Aspergillus glaucus</i> <i>Scopulariopsis brevicaulis</i> <i>Aspergillus fumigatus</i>  <i>Aureobasidium pullulans</i>	15,100 27,800 69,400 41,700 13,900 <b>Synthetic pillows:</b> 2745 <b>Feather pillows:</b> 1863 <b>Synthetic pillows:</b> 1926 <b>Feather pillows:</b> 5110	CFU/g pillow dust Mean  CFU/g pillow material (not dust) Mean	Not reported
20. Krämer et al. (2006) [80]	Relationship between eczema and exposure to HDM allergens	Augsburg, Germany	Der p 1 Der f 1	0.68 (1.08, 16.0) 0.79 (1.3, 22.0)	µg/g mattress dust Mean (Median, Maximum)	Not reported
21. Chen et al. (2007) [26]	Explored associations between socioeconomic status and the concentrations of indoor bio-contaminants	Munich & Leipzig, Germany	Der p 1  Der f 1	<b>Mothers' Mattress:</b> 97.7 (5–608.3) <b>Infants' Mattress:</b> 5.0 (5.0–642.9) <b>Mothers' Mattress:</b> 634.1 (125.4–3601.8) <b>Infants' Mattress:</b> 525.1 (5.0–2324.7)	ng/g mattress dust (Median (25 <sup>th</sup> –75th percentiles))  ng/g mattress dust (Median (25 <sup>th</sup> –75th percentiles))	Mothers' Mattress:  0.7385 (0.453–1.166)

(continued on next page)



Table 1 (continued)

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
22. Krop et al. (2007) [82]	Assessed the presence of occupational laboratory animal allergens in mattress dust of workers	Netherlands	Fel d 1	<b>Mothers' Mattress:</b> 180.2 (68.6–820.5) <b>Infants' Mattress:</b> 326.0 (84.7–1322.5)	ng/g mattress dust (Median (25 <sup>th</sup> –75th percentiles))	Infants' Mattress:
			Endotoxin	<b>Mothers' Mattress:</b> 3008 (1045.5–7913) <b>Infants' Mattress:</b> 5866 (2336–14,669)	EU/g mattress dust (Median (25 <sup>th</sup> –75th percentiles))	0.188 (0.120–0.289)
			Rat urinary allergen	<b>Laboratory animal workers:</b> 39.3 (19.8–78.0) <b>Non-laboratory animal workers (controls):</b> 7.6 (4.7–12.2)	ng/g mattress dust (Median (Range))	Not reported
			Mouse urinary allergen	<b>Laboratory animal workers:</b> 29.5 (11.7–74.6) <b>Non-laboratory animal workers (controls):</b> 8.8 (4.6–16.8)		
23. Giovannangelo et al. (2007) [54]	Measured levels of $\beta$ -Glucans and fungal extracellular polysaccharide in mattress dust of children	Germany, Netherlands & Sweden	(1–3)- $\beta$ -Glucan	<b>Germany:</b> 450 (150–1050) (level), 1050 (900–2050) (load) <b>Netherlands:</b> 400 (150–1000) (level), 1050 (750–3000) (load) <b>Sweden:</b> 350 (100–1000) (level), 225 (1050–4000) (load)	Level: $\mu$ g/m <sup>2</sup> mattress area sampled Load: $\mu$ g/g mattress dust (–Median (95% CI))	Germany: ~0.25 (0.01–1) Netherlands: ~0.25 (0.01–1) Sweden: ~0.15 (0.02–0.9) (–Median (Range))
			Fungal Extracellular Polysaccharide (EPS)	<b>Germany:</b> 10,000 (2000–50,000) (level), 40,000 (10,500–100,000) (load) <b>Netherlands:</b> 9000 (1050–45,000) (level), 40,000 (10,500–100,000) (load) <b>Sweden:</b> 2050 (400–10,500) (level), 15,000 (4000–50,000) (load)	Level: EPSU/m <sup>2</sup> mattress area sampled Load: EPSU/g mattress dust (–Median (95% CI))	
24. Rennie et al. (2008) [125]	Investigated associations between endotoxin levels and asthma and wheeze in children	Saskatchewan, Canada	Endotoxin	2069.79 (4650.02) (level), 7.99 (9.85) (load)	Level: EU/m <sup>2</sup> mattress area sampled Load: EU/mg mattress dust (Median (IQR))	0.365 $\pm$ 0.26 (Mean $\pm$ SD)
25. Nam et al. (2008) [110]	Measured HDM allergen concentrations in house dust	Cheonan, Korea	Der f 1	4.0 (0.01–112.1) (level), 10.2 (0.01–230.9) (load)	Level: $\mu$ g/m <sup>2</sup> mattress area sampled	Not reported
			Der p 1	0.02 (0.01–8.8) (level), 0.14 (0.01–30.0) (load)	Load: $\mu$ g/g mattress dust (–Median (Range))	
26. Korhals et al. (2008) [79]	Investigated the occurrence of <i>Listeria</i> spp. bacteria in mattress dust of farm children	Bavaria, Germany	<i>Listeria</i> spp. <i>L. monocytogenes</i>	Detected in 8% of mattress dust samples (Culture methods) Detected in 3% of mattress dust samples (Culture methods) Detected in 8% of mattress dust samples (Real-time PCR methods) Detected in 8% of mattress dust samples (Culture methods)		Not reported
27. Vogel et al. (2008) [164]	Identified microbial flora on the mattresses of farm children	Bavaria, Germany	<i>L. innocua</i> <i>Bacillus</i> spp. Coliform bacteria <i>Lactobacillus</i> spp. <i>Enterococcus</i> spp. <i>Clostridium</i> spp. <i>Aspergillus</i> spp. <i>Mucor</i> spp. <i>Cladosporium</i> spp.	5.25 (0.87) 5.67 (0.94) 3.00 (3.30) 3.15 (4.47) 0.00 (3.00) 4.00 (1.10) 3.75 (1.00) 0.00 (3.74)	log. CFU (Median (IQR))	Not reported

			<i>Penicillium</i> spp.	3.00 (4.00)		
			<i>Wallemia</i> spp.	3.48 (4.00)		
28. Soleimani and Rafinejad (2008) [139]	Analyzed HDM contamination in hotels and inns	Bandar-Abbas, Iran	<b>HDM species:</b> <i>D. pteronyssinus</i> (identified in 84.5% of samples), <i>D. farinae</i> (identified in 13% of samples), <i>C. malaccensis</i> (identified in 2.5% of samples)			Not reported
29. Heyes et al. (2009) [62]	Measured (1–3)- $\beta$ -Glucan levels in pillows, duvets, and mattresses	Wellington, New Zealand	(1–3)- $\beta$ -Glucan	<b>Mattress:</b> 15.7 (10.4–23.9) (level), 76.6 (61.4–94.0) (load) <b>Duvets:</b> 8.8 (5.4–14.1) (level), 132.1 (68.9–207.9) (load) <b>Pillows:</b> 3.5 (2.5–4.8) (level, $\mu\text{g}$ pillow), 132.1 (68.9–207.9) (load) <b>Synthetic Pillows:</b> 3.8 (1.4–11.0) (level, $\mu\text{g}$ /pillow), 105.4 (76.9–144.6) (load) <b>Feather Pillows:</b> 2.0 (0.8–5.1) (level, $\mu\text{g}$ /pillow), 76.6 (36.7–159.9) (load)	Level: $\mu\text{g}/\text{m}^2$ mattress/pillow/duvet area sampled Load: $\mu\text{g}/\text{g}$ mattress/pillow/duvet dust (Mean (95% CI))	Not reported
30. Wu et al. (2009) [172]	Measured allergen and microbial bio-contaminant concentrations in homes of asthmatic children	Central Taiwan	Der p 1	<b>Pillow:</b> 1.33 (0.91–1.94) <b>Bamboo side of mattress:</b> 0.38 (0.25–0.58) <b>Conventional side of mattress:</b> 0.77 (0.51–1.17)	$\mu\text{g}/\text{g}$ mattress/pillow dust (Mean (95% CI))	Not reported
			Der f 1	<b>Pillow:</b> 0.22 (0.14–0.33) <b>Bamboo side of mattress:</b> 0.24 (0.14–0.44) <b>Conventional side of mattress:</b> 0.55 (0.30–1.01)		
			Blo t 5	<b>Pillow:</b> 0.014 (0.011–0.017) <b>Bamboo side of mattress:</b> 0.024 (0.017–0.033) <b>Conventional side of mattress:</b> 0.165 (0.115–0.237)		
			Total HDM allergen	<b>Pillow:</b> 2.72 (1.99–3.73) <b>Bamboo side of mattress:</b> 1.35 (0.95–1.91) <b>Conventional side of mattress:</b> 3.50 (2.65–4.62)		
			Endotoxin	<b>Bamboo side of mattress:</b> 163.7 (82.6–244.8) <b>Conventional side of mattress:</b> 358.2 (211.7–504.8)	EU/mg mattress dust (Mean (95% CI))	
			$\beta$ -Glucan	<b>Bamboo side of mattress:</b> 27.9 (22.4–34.8) <b>Conventional side of mattress:</b> 25.5 (21.9–29.8)	$\mu\text{g}/\text{g}$ mattress dust (Mean (95% CI))	
31. Chen et al. (2009) [27]	Investigated associations between infant exposure to endotoxins and development of eczema	Munich & Leipzig, Germany	Endotoxin	<b>Mothers' mattress:</b> 2071 (595–6919) (level), 3008 (1046–7913) (load) <b>Children's mattress:</b> 1015 (330–3022) (level), 5866 (2336–14,669) (load)	Level: EU/m <sup>2</sup> mattress area sampled Load: EU/g mattress dust (Median (IQR))	Not reported
32. Täubel et al. (2009) [151]	Explored the origins of bacteria in house dust	Finland	Bacterial phyla and genera	<b>Gram-positive bacterial groups (phylum:genera) identified in mattress dust:</b> Actinobacteria: <i>Corynebacterium</i> , <i>Propionibacterium</i> , <i>Micrococcus</i> Firmicutes: <i>Staphylococcus</i> , <i>Lactobacillus</i> , <i>Streptococcus</i> , <i>Lactococcus</i> Deinococcus-Thermus: <i>Deinococcus</i> <b>Gram-negative bacterial phyla identified in mattress dust:</b> Proteobacteria: <i>Paracoccus</i> , <i>Aggregatibacter</i> Bacteroidetes: <i>Prevotella</i> , <i>Bacteroides</i>		Not reported

(continued on next page)

Table 1 (continued)

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
			Endotoxin	1.4 ± 3.7 (load), 194.1 ± 5.8 (level)	Load: EU/mg mattress dust Level: EU/m <sup>2</sup> mattress area sampled (Mean ± SD)	
			Lipopolysaccharide (LPS)	0.031 ± 1.379 (load), 4.396 ± 1.944 (level)	Load: nmol/mg mattress dust Level: nmol/m <sup>2</sup> mattress area sampled (Mean ± SD)	
			Muramic acid	15.5 ± 1.7 (load), 2191.2 ± 3.3 (level)	Load: ng/mg mattress dust Level: ng/m <sup>2</sup> mattress area sampled (Mean ± SD)	
33. Eberlein et al. (2009) [42]	Measured endotoxin and allergen levels in hospitals and adjacent homes	Bavaria, Germany	Endotoxin	<b>Hospitals:</b> 321,720 (230,500–518,700) <b>Adjacent Homes:</b> 243,980 (156,520–399,020)	EU/g mattress dust (Median (IQR))	Not reported
			Der p 1	<b>Hospitals:</b> 0 (0–208) <b>Adjacent Homes:</b> 163 (0–391)	ng/g mattress dust (Median (IQR))	
			Der f 1	<b>Hospitals:</b> 409 (152–839) <b>Adjacent Homes:</b> 791 (320–2074)		
			Fel d 1	<b>Hospitals:</b> 663 (117–2426) <b>Adjacent Homes:</b> 1227 (261–2824)		
34. Spertini et al. (2009) [140]	Evaluation of the reduction in allergens in houses with improved energy performance	Lausanne, Switzerland	Der f 1	<b>Control Buildings (CB):</b> 954; <b>Low Energy Use Buildings (LEB):</b> 67 <b>CB:</b> 73; <b>LEB:</b> 51 <b>CB:</b> 15; <b>LEB:</b> 5	ng/g mattress dust (Median)	Not reported
35. Visitsunthorn et al. (2010) [163]	Measured HDM allergen concentrations in different types of mattresses	Bangkok, Thailand	Der p 1 Mite Group 2 Der p 1 + Der f 1	<b>Coconut mattress:</b> 0.96 (0.39–1.45) (month 1), 16.60 (0–33.7) (month 12) <b>Kapok mattress:</b> 1.44 (1.06–1.83) (month 1), 31.30 (17.40–42.80) (month 12) <b>Sponge (polyurethane foam) mattress:</b> 0.96 (0.23–1.75) (month 1), 12.90 (1.00–17.90) (month 12) <b>Synthetic mattress:</b> 1.30 (0.80–1.89) (month 1), 25.30 (12.10–32.50) (month 12)	µg/g mattress dust (Median (IQR))	Not reported
36. Doyen et al. (2011) [40]	Measured endotoxin concentrations in mattresses at time of birth of newborn and at 6 months of life	Belgium	Endotoxin	<b>At birth, mattresses in newborn's bedroom:</b> 17.6 (0.4–346.7) (Site A), 20.4 (0.8–26.3) (Site B) <b>6 months of life, mattresses in newborn's bedroom:</b> 79.6 (3.8–518.8) (Site A), 101.8 (6.5–634.4) (Site B)	EU/mg mattress dust (Median (Range))	Not reported
37. Leung et al. (2011) [93]	Measured allergen levels in settled dust in residences of asthmatic children	Hong Kong	Fel d 1	0.06 (0.03–0.10)	µg/g mattress dust (Median (IQR))	Not reported
			Bla g 2	24.8 (14.2–47.7)	ng/g mattress dust (Median (IQR))	
			Blo t 5	0.26 (0.14–0.79)	µg/g mattress dust (Median (IQR))	

38. Tischer et al. (2011) [153]	Investigated exposure of children to endotoxins, $\beta$ -Glucans, and EPS mold components in house dust	Germany & Netherlands	Endotoxin  (1–3)- $\beta$ -D-Glucan  Extracellular polysaccharides (EPS)	<b>German LISA &amp; GINI Studies:</b> 3053 (1521–6015) (level), 12,222 (7379–21,337) (load) <b>Dutch PIAMA Study:</b> 2356 (1461–4208) (level), 10,608 (6550–17,366) (load) <b>German LISA &amp; GINI Studies:</b> 421 (238–865) (level), 1859 (1277–2396) (load) <b>Dutch PIAMA Study:</b> 380 (199–625) (level), 1662 (1135–2205) (load) <b>German LISA &amp; GINI Studies:</b> 1008 (4458–25,904) (level), 40,792 (24,235–65,371) (load) <b>Dutch PIAMA Study:</b> 8257 (3890–17,310) (level), 34,696 (20,364–58,156) (load)	Level: EU/m <sup>2</sup> mattress area sampled Load: EU/g mattress dust (Median (IQR)) Level: $\mu$ g/m <sup>2</sup> mattress area sampled Load: $\mu$ g/g mattress dust (Median (IQR)) Level: EPSU/m <sup>2</sup> mattress area sampled Load: EPSU/g mattress dust (Median (IQR))	German LISA & GINI Studies: 0.257 (0.139–0.471) Dutch PIAMA Study: 0.247 (0.148–0.366) (Median (IQR))
39. Hsu et al. (2012) [68]	Explored associations between indoor phthalate concentrations and mold growth	Tainan, Taiwan	DEHP  BBzP  DBP	<b>w/o visible mold:</b> 656.6 (459.4–980.9) <b>w/ visible mold:</b> 933.3 (580.0–1404.9) <b>w/o visible mold:</b> 2.1 (1.0–6.4) <b>w/ visible mold:</b> 3.7 (1.0–6.1) <b>w/o visible mold:</b> 16.6 (10.4–22.8) <b>w/ visible mold:</b> 20.2 (12.4–40.9)	$\mu$ g/g mattress dust (Median (25 <sup>th</sup> –75 <sup>th</sup> percentiles))	Not reported
40. Ege et al. (2012) [45]	Identified environmental bacteria in dust on children's mattresses	Bavaria, Germany		<b>Bacterium identified which are related to a physician's diagnosis of asthma:</b> <i>Corynebacterium mycetoides</i> , <i>Zoogloea</i> sp., <i>Duganella</i> sp., <i>Aurantimonas ureolytica</i> sp., <i>Serratia marcescens</i> , <i>S. nematodiphila</i> , <i>Serratia</i> sp., <i>Pseudomonas uorescens</i> , <i>Corynebacterium tuberculostearicum</i> , <i>Gardnerella vaginalis</i> , <i>Lactobacillus curvatus</i> , <i>Lactobacillus sakei</i> , <i>Streptococcus</i> sp., <i>Moraxella</i> sp., <i>Staphylococcus sciuri</i> sp., <i>Jeotgalicoccus</i> sp., <i>Salinicoccus</i> sp., <i>Macrococcus brunensis</i> , <i>Bacillus</i> sp. <b>Bacterium identified which are related to hay fever:</b> <i>Aerobacter ureolytica</i> , <i>P. fluorescens</i> , <i>Corynebacterium mucifaciens</i> , <i>C. freiburgense</i> , <i>C. variabile</i> , <i>C. sp.</i> , <i>Triatoma infestans</i> , <i>Neisseria meningitidis</i> , <i>N. mucosa</i> , <i>N. subflava</i> <b>Bacterium identified which are related to atopic sensitization:</b> <i>Enterobacter cloacae</i> , <i>E. aerogenes</i> , <i>E. cancerogenus</i> , <i>E. ludwigii</i> , <i>Pantoea</i> sp., <i>Kluyvera cryocrescens</i> , <i>Erwinia persicina</i> , <i>Aurantimonas altamirensis</i> ; <i>A. ureolytica</i> , <i>Mesorhizobium</i> sp., <i>Lactobacillus iners</i> , <i>Acinetobacter lwoffii</i>		Not reported
41. Barnig et al. (2012) [5]	Measured endotoxin concentrations in the dust of rural and urban residences	Strasbourg & Haut-Doubs, France	Endotoxin	<b>Type of farm:</b> Non-modern: 3700 $\pm$ 4900 Modern: 1200 $\pm$ 1600 <b>Barn position:</b> Adjacent to house: 3400 $\pm$ 4700 Separated to house: 2200 $\pm$ 3600 <b>Storage hay mode:</b> Round bales: 3200 $\pm$ 4700 In bulk: 1100 $\pm$ 900 Other: 2700 $\pm$ 3600	ng/g mattress dust (Mean)	Not reported
42. Begum et al. (2012) [7]	Evaluated the prevalence of fungi in used pillows	Perth, Australia	Fungi: <i>Alternaria</i> sp., <i>Aureobasidium</i> sp., <i>Cladosporium</i> sp., <i>Epicoccum</i> sp., <i>Monilia</i> sp., <i>Penicillium</i> sp., Yeast, <i>Rhizopus</i> sp.	<b>Total fungal counts:</b> 241 (22–563) (Yeast most prevalent)	CFU/pillow Mean (Range)	0.06–0.36 g pillow dust/pillow
43. Lawson et al. (2012) [91]	Measured endotoxin concentrations and loading in mattress dust	Humboldt, Saskatchewan, Canada	Endotoxin	<b>Children with asthma or wheeze (cases):</b> 19.6 (15.6–24.8)	EU/mg mattress dust (Mean (95% CI))	Not reported

(continued on next page)

Table 1 (continued)

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
44. Ali et al. (2012) [2]	Identified and quantified flame retardants in floor and mattress dust	Wellington, Wairarapa, Christchurch, & North Canterbury, New Zealand	OPFRs  Not reported	<b>Children without asthma or wheeze (controls):</b> 21.1 (15.6–24.8) <b>Children with asthma or wheeze (cases):</b> 240.5 (212.1–370.9) <b>Children without asthma or wheeze (controls):</b> 376.2 (324.4–436.2)	EU/m <sup>2</sup> mattress area sampled (Mean (95% CI)) TEP: 10 (<10–235), TnBP: 65 (20–1920), TCEP: 38 (<20–475), TCPP: 250 (133–1920), TBEP: 1545 (635–3610), TPhP: 240 (20–35,190), TDCPP: 103 (20–6465), TCP: 157 (<50–2155) $\alpha$ -HBCD: 41 (3–345), $\beta$ -HBCD: 9 (<2–122), $\gamma$ -HBCD: 80 (18–1260) BTBPE: 1 (<2–37), TBB: 3 (<2–40), TBPH: 1 (<2–50), DBDPE: 9 (<5–220)	ng/g mattress dust (Median (Range))
45. Wu et al. (2012) [173]	Assessed the impact of daily vacuuming of mattresses to reduce levels of HDM allergens, bacterial endotoxin and fungal $\beta$ -Glucans	Taiwan	Der p 1 Der f 1 Endotoxin $\beta$ -Glucan	<b>Week 0:</b> 0.68 (0.32–1.40), <b>Week 8:</b> 0.46 (0.23–0.92) <b>Week 0:</b> 1.18 (0.60–2.30), <b>Week 8:</b> 0.69 (0.41–1.14) <b>Week 0:</b> 91 (64–130), <b>Week 8:</b> 97 (75–125) <b>Week 0:</b> 63.2 (46.1–86.6), <b>Week 8:</b> 75.1 (54.1–104.0)	$\mu$ g/g mattress dust (Mean (95% CI)) EU/mg mattress dust (Mean (95% CI)) $\mu$ g/g mattress dust (Mean (95% CI))	1.5 (1.12–2) at Week 0, to 0.27 (0.18–0.41) at Week 8 (Mean (95% CI))
46. Gehring et al. (2012) [53]	Assessed the impact of mite-impermeable mattress covers on allergen exposure	Netherlands	Der f 1 Der p 1	<b>Placebo mattress cover:</b> 2849 $\pm$ 5.7 <b>Active mattress cover:</b> 886 $\pm$ 3.5 <b>Placebo mattress cover:</b> 410 $\pm$ 4.7 <b>Active mattress cover:</b> 397 $\pm$ 3.5	ng/g mattress dust (Mean $\pm$ SD)	Not reported
47. Casas et al. (2013) [23]	Analyzed the results of the European HITEA project (Health Effects of Indoor Pollutants: Integrating microbial, toxicological, and epidemiological approaches) with 4 European birth cohorts	Finland, Germany, Netherlands, & Spain	Endotoxin  Fungal Extracellular Polysaccharide (EPS)  (1–3)- $\beta$ -Glucan	<b>German LISA Study:</b> 23.0 (21.1–25.1) <b>Dutch PIAMA Study:</b> 22.0 (19.8–24.5) <b>Spanish INMA Study:</b> 3.2 (2.7–3.9) <b>Finnish LUKAS2 Study:</b> 17.6 (15.8–19.5) <b>German LISA Study:</b> 47.5 (43.7–51.6) <b>Dutch PIAMA Study:</b> 30.7 (27.8–33.8) <b>Spanish INMA Study:</b> 116.7 (107.2–127.0) <b>Finnish LUKAS2 Study:</b> 39.6 (34.1–46.0) <b>German LISA Study:</b> 0.9 (1.8–2.0) <b>Dutch PIAMA Study:</b> 1.7 (1.5–1.8)	EU/mg mattress dust (Mean (95% CI)) EPSU/mg mattress dust (Mean (95% CI)) $\mu$ g/mg mattress dust (Mean (95% CI))	Not reported

48. Callesen et al. (2014) [21]	Evaluated associations between selected allergens and asthma, rhinoconjunctivitis, and atopic dermatitis in preschool children	Odense, Denmark	Der p 1	<b>Spanish INMA Study:</b> - <b>Finnish LUKAS2 Study:</b> 2.4 (2.2–2.7) <b>Healthy Controls:</b> 110.0 <b>Asthma:</b> 132.2, <b>s-IgE</b> -: 145.4, <b>s-IgE</b> + : 132.0 <b>Rhinoconjunctivitis:</b> 119.4, <b>s-IgE</b> -: 164.9, <b>s-IgE</b> + : 104.8 <b>Atopic dermatitis:</b> 131.8, <b>s-IgE</b> -: 176.0, <b>s-IgE</b> + : 151.0	ng/g mattress dust (Median)	Not reported
			Der f 1	<b>Healthy Controls:</b> 977.2 <b>Asthma:</b> 988.4, <b>s-IgE</b> -: 471.8, <b>s-IgE</b> + : 1726 <b>Rhinoconjunctivitis:</b> 1,251, <b>s-IgE</b> -: 173.7, <b>s-IgE</b> + : 988.4 <b>Atopic dermatitis:</b> 751.9, <b>s-IgE</b> -: 672.4, <b>s-IgE</b> + : 1211		
			Der p 2	<b>Healthy Controls:</b> 121.5 <b>Asthma:</b> 119.5, <b>s-IgE</b> -: 69.2, <b>s-IgE</b> + : 135.5 <b>Rhinoconjunctivitis:</b> 153.4, <b>s-IgE</b> -: 158.1, <b>s-IgE</b> + : 130.4 <b>Atopic dermatitis:</b> 119.5, <b>s-IgE</b> -: 88.3, <b>s-IgE</b> + : 130.0		
			Fel d 1	<b>Healthy Controls:</b> 114.5 <b>Asthma:</b> 76.7, <b>s-IgE</b> -: 210.7, <b>s-IgE</b> + : 42.1 <b>Rhinoconjunctivitis:</b> 85.6, <b>s-IgE</b> -: 61.6, <b>s-IgE</b> + : 99.6 <b>Atopic dermatitis:</b> 150.4, <b>s-IgE</b> -: 170.9, <b>s-IgE</b> + : 116.6		
			Can f 1	<b>Healthy Controls:</b> 81.6 <b>Asthma:</b> 71.1, <b>s-IgE</b> -: 77.7, <b>s-IgE</b> + : 71.8 <b>Rhinoconjunctivitis:</b> 54.2, <b>s-IgE</b> -: 28.5, <b>s-IgE</b> + : 66.4 <b>Atopic dermatitis:</b> 53.0, <b>-IgE</b> -: 53.3, <b>s-IgE</b> + : 57.2		
49. Valkonen et al. (2015) [160]	Evaluated associations between bacterial exposures and atopy and asthma in children	Germany, Austria, and Switzerland	<i>Mycobacterium</i> spp.	<b>Asthma:</b> Farm Children (FC): 1,252,815, Exposed Non-Farm Children (ENFC): 738,266, Non-Exposed Non-Farm Children (NENFC): 326,199 <b>No asthma, but atopy:</b> FC: 872,049, ENFC: 583,315, NENFC: 360,554 <b>No asthma or atopy:</b> FC: 868,964, ENFC: 876,830, NENFC: 690,282	cells/m <sup>2</sup> mattress area sampled (Median)	Not reported
			<i>Bifidobacteriaceae</i> spp.	<b>Asthma:</b> FC: 7,853,091, ENFC: 4,933,515; NENFC: 4,337,218 <b>No asthma, but atopy:</b> FC: 6,882,662; ENFC: 2,054,042; NENFC: 3,317,576 <b>No asthma or atopy:</b> FC: 5,168,782; ENFC: 4,330,676, NENFC: 6,571,929		

(continued on next page)

Table 1 (continued)

Study	Description	Location of Field Study	Detected Biological Material or Chemical Contaminant	Mattress Dust Concentration	Units	Mattress Dust Load, $m_0$ (g/m <sup>2</sup> )
			<i>Clostridium</i> cluster I	<b>Asthma:</b> FC: 264,510; ENFC: 32,028; NENFC: 31,438 <b>No asthma, but atopy:</b> FC: 131,510; ENFC: 30,665; NENFC: 14,547 <b>No asthma or atopy:</b> FC: 106,416; ENFC: 32,147; NENFC: 73,044		
			<i>Clostridium</i> cluster XI	<b>Asthma:</b> FC: 277,263; ENFC: 54,883; NENFC: 54,362 <b>No asthma, but atopy:</b> FC: 172,193; ENFC: 24,904; NENFC: 35,117 <b>No asthma or atopy:</b> FC: 271,469; ENFC: 54,771; NENFC: 90,370		
50. Tischer et al. (2015) [154]	Evaluated associations between bacteria and fungi concentrations and asthma and lung function	Iceland (Reykjavik), Sweden (Uppsala, Umea, Gothenburg), Estonia (Tartu), Germany (Hamburg, Erfurt), UK (Norwich, Ipswich), Belgium (Antwerp Centre, Antwerp South), France (Paris, Grenoble), Switzerland (Basel), Italy (Verona, Pavia, Turin), and Spain (Oviedo, Galdakao, Barcelona, Albacete, Huelva)	<i>Cladosporium herbarum</i> <i>Penicillium</i> spp./ <i>Aspergillus</i> spp./ <i>Paecilomyces variotii</i> group (1–3)- $\beta$ -Glucan <i>Mycobacterium</i> spp. Muramic acid Gram-Positive bacteria Gram-Negative bacteria	64 (2; 3776) 39,340 (638; 261,600,000) 0.87 (0.08; 7) 5413 (258; 524,700) 15 (1; 116) 568,400 (16,480; 65,870,000) 66,070 (1945; 6,900,000)	cells/mg mattress dust (Median (Minimum; Maximum)) $\mu$ g/mg mattress dust (Median (Minimum; Maximum)) cells/mg mattress dust (Median (Minimum; Maximum)) $\mu$ g/mg mattress dust (Median (Minimum; Maximum)) cells/mg mattress dust (Median (Minimum; Maximum))	Not reported
51. Canbaz et al. (2016) [22]	Evaluated associations between organophosphate and polybrominated diphenyl ether flame retardants in dust and asthma	Sweden	OPFRs  BFRs	<b>Asthmatic children:</b> TCEP: 102 (49–263), TCPP: 116 (60–243), TDCIPP: 141 (85–427), TBOEP: 3138 (1082–5593), TPhP: 163 (107–276), EHDPHP: 163 (107–276), mmp-TMPP: 192 (90–437), $\Sigma$ OPFRs: 6449 (3718–12,865) <b>Control group:</b> TCEP: 107 (52–207), TCPP: 96 (44–362), TDCIPP: 162 (88–374), TBOEP: 3723 (1434–7069), TPhP: 631 (308–1493), EHDPHP: 172 (111–421), mmp-TMPP: 288 (102–790), $\Sigma$ OPFRs: 6795 (3575–13,000) <b>Asthmatic children:</b> BDE-47: 11 (6–24), BDE-99: 14 (7–36), BDE-100: 2 (1–5), BDE-153: 7 (3–20), BDE-183: 3 (1–9), BDE-209: 118 (49–306), $\Sigma$ BFRs: 259 (107–498) <b>Control group:</b> BDE-47: 11 (7–24), BDE-99: 13 (8–34), BDE-100: 3 (2–6), BDE-153: 6 (3–16), BDE-183: 3 (1–10), BDE-209: 122 (36–389), $\Sigma$ BFRs: 239 (91–7910)	ng/g mattress dust (Median (25 <sup>th</sup> –75 <sup>th</sup> percentiles))	Not reported

52. Stein et al. (2016) [146] Assessment of asthma risk in Amish and Hutterite farm children Indiana and South Dakota, USA Not reported

**Bacterial families with a relative abundance  $\geq 2\%$  in Amish or Hutterite dust, relative proportion in each group shown in parentheses:**

Corynebacteriaceae (0.026 vs. 0.067), Micrococaceae (0.045 vs. 0.04), Prevotellaceae (0.001 vs. 0.022), Bacillaceae (0.041 vs. 0.021), Staphylococaceae (0.037 vs. 0.08), Lactobacillaceae (0.003 vs. 0.02), Streptococaceae (0.022 vs. 0.065), Clostridiaceae (0.007 vs. 0.046), Lachnospiraceae (0.005 vs. 0.026), Ruminococaceae (0.014 vs. 0.047), Bartonellaceae (0.347 vs. 0), Rhodobacteraceae (0.025 vs. 0.016), Sphingomonadaceae (0.033 vs. 0.042), Enterobacteriaceae (0.029 vs. 0.021), Moraxellaceae (0.032 vs. 0.026)

**Nomenclature:** HDM: House Dust Mite; CFU: Colony Forming Units; EU: Endotoxin Units; EPSU: Fungal Extracellular Polysaccharide (EPS) Units; PCR: Polymerase Chain Reaction; IgE: Immunoglobulin E antibody; CI: Confidence Interval; IQR: Interquartile Range; ND: Not Detected; DEHP: Bis(2-ethylhexyl) phthalate; BBzP: Benzyl butyl phthalate; DBP: Dibutyl phthalate; OPFRs: Organophosphate flame retardants; TEP: tri-ethyl-phosphate; TnBP: tri-*n*-butyl-phosphate; TCEP: tris-(2-chloroethyl)-phosphate; TCEP: tris-(2-chloroethyl)-phosphate; TCPP: tris-(1-chloro-2-propyl) phosphate; TDCCP: tris-(2,3-dichloropropyl)-phosphate; TDCCPP: tris-(1,3-dichloroisopropyl) phosphate; TBOEP: tris(2-butoxyethyl) phosphate; TPhP: tri-phenyl-phosphate; EHDHP: 2-ethylhexyldiphenyl phosphate; mmp-TMPP: meta, meta, para-tricresylphosphate; BFRs: Brominated flame retardants; HBCDs: Hexabromocyclododecanes;  $\alpha$ -HBCD,  $\beta$ -HBCD,  $\gamma$ -HBCD; NBFRs: Novel Brominated flame retardants; BTBPE: 1,2-bis(2,4,6-tribromophenoxy)ethane; DBDPE: decabromodiphenyl ethane; TBB: 2-ethylhexyl-2,3,4,5-tetrabromobenzoate; TBPH: bis(2-ethylhexyl)-3,4,5,6-tetrabromophthalate; Der p 1: Dermatophagoides pteronyssinus Group 1 (European HDM); Der p 2: Dermatophagoides pteronyssinus Group 2 (European HDM); Der f 1: Dermatophagoides farinae Group 1 (American HDM); Can f 1: Canis familiaris Group 1 (Dog allergen); Fel d 1: Felis domesticus Group 1 (Cat allergen); Phi p 5: Phleum pretense Group 5 (Common timothy pollen); Bla g 2: Aspartic protease Group 2 (Cockroach allergen); Bio t 5: Blomia tropicalis Group 5 (Blomia tropicalis HDM); LISA Study: Lifestyle Related Factors on the Immune System and the Development of Allergies in Childhood (Germany); GINI Study: German Infant Nutritional Intervention (Germany); PIAMA Study: Prevention and Incidence of Asthma and Mite Allergy (Netherlands); INMA Study: Infancia y Medio Ambiente (Environment and Childhood) (Spain); LUKAS2 Study: Lapsuuden kasvuympäristö ja allergiat 2 (Childhood environment and allergies) (Finland).

marker used to assess fungal exposure. Typical concentrations are in the range of  $10\text{--}10^3$   $\mu\text{g/g}$  of mattress dust. Mattresses and pillows are an ideal fungal culture medium, given the high moisture levels induced by human sweating ( $\sim 100$  L of sweat per year) and elevated temperatures ( $>30$  °C) [36,171]. Fungi may consume HDM fecal matter and squames in mattress dust, providing them with necessary nutrients; and subsequently, HDMs may consume fungi, leading to a “miniature ecosystem” within mattresses and pillows, as suggested by Refs. [78,171].

Similar fungal species have been detected in mattress dust throughout the world, including the United Kingdom [171], Northern California, U.S. [63], Germany [73,119,164,200], and Australia [7]. Refs. [23,54,153] found small differences in (1–3)- $\beta$ -Glucan levels in mattress dust in large cohort studies across several European countries. Additionally, the fungal composition of mattress dust can vary with bedding material, as observed by Ref. [119] for dust on cotton and synthetic mattress encasings. Higher concentrations of *Aspergillus fumigatus* and (1–3)- $\beta$ -Glucans have been reported in synthetic compared to feather pillows [171,62].

### 3.3. Bacteria

The sleep microenvironment is also home to an array of bacterial phyla and genera, many of which are associated with human origins (skin, oral, intestinal/fecal, and genital) and specifically, the shedding of human skin, e.g. Refs. [45,66,151,154]. Common bacterium identified in mattress dust include: *Staphylococcus*, *Lactobacillus*, *Streptococcus* sp., *Lactococcus*, *Bacillus* sp., *Listeria* spp., *Zoogloea* sp., *Duganella* sp., *Aurantimonas ureolytica* sp., *Serratia marcescens*, *Pseudomonas fluorescens*, *Corynebacterium tuberculostearicum*, *Gardnerella vaginalis*, *Lactobacillus curvatus*, *Moraxella* sp., *Staphylococcus sciuri* sp., *Macroccoccus brunensis*, *Aerobacter ureolytica*, *P. fluorescens*, *Triatoma infestans*, and *Neisseria meningitidis*, among a host of others. Concentrations can range from  $< 10^6$  to  $> 10^8$  cells per g of mattress dust (Table 1). Differences in the bacterial composition of mattress dust have been observed among Amish and Hutterite families in the United States [146] and among children from farming and non-farming households in Europe [160]. In addition to identifying specific bacterium, endotoxin concentrations are also commonly reported in mattress dust. Endotoxin is the biologically active lipopolysaccharide (LPS) of gram-negative bacteria and is a common marker to assess bacterial exposure (although it only represents defined subpopulations of bacteria and does not differentiate between human and non-human sources) [151] and may also serve as an indicator of inflammatory potential of settled dust [100]. Endotoxin levels are reported as endotoxin units (EU) per gram of mattress dust, and concentrations are typically in the range of  $10^3$  to  $10^5$  EU/g. Greater levels of bacterial endotoxin loading have been found in infant mattresses, compared to those of their mothers [26,52]. Furthermore, Doyen et al. [40] found endotoxin levels to increase in infants' bedrooms over the first 6 months of life. As with fungal (1–3)- $\beta$ -Glucans, small variability in endotoxin levels have been found among large cohort studies in Europe [23,153].

### 3.4. Plasticizers and flame retardants

Due to their low volatility and high molecular weight, SVOCs, such as plasticizers and flame retardants, tend to accumulate in house dust, e.g. Refs. [69,167]. Although many field studies have measured the concentrations of various SVOCs in settled dust on flooring, only a few have explicitly reported levels in mattress dust. [68] detected several phthalate plasticizers in mattress dust in Taiwan, including DEHP, BBzP, and DBP (nomenclature listed in



**Table 1**). Concentrations are on the order of  $10\text{--}10^2$   $\mu\text{g/g}$  of mattress dust, with higher levels reported in the presence of visible mold on the mattress. [2,22] identified numerous flame retardants in mattress dust, including polybrominated diphenyl ethers (e.g. BDE 47, BDE 99, BDE 183, BDE 197, and BDE 209), organophosphates (e.g. TEP, TCEP, TCPP, and TDCPP, among others), HBCDs, and novel flame retardants (BTBPE, TBB, and TBH, among others). Concentrations for the various flame retardants ranged over several orders of magnitude, from  $<1$  to  $10^3$  ng/g mattress dust. Plasticizers and flame retardants may be found as chemical additives in mattresses, especially infant crib mattresses [17,32,144]. It is likely that some of these compounds partitioned directly into settled mattress dust as they migrated from the underlying mattress foam and encasing, whereas others may have originated from common indoor sources, such as vinyl flooring, foam furniture, and electronic casings, and then deposited (after partitioning to airborne particles) on the mattress and bedding.

### 3.5. Particle deposits on mattresses and bedding materials

The mass of dust that accumulates on mattresses, pillows, and bedding is an important parameter that can affect exposure and surface-to-air transport of the settled particles, e.g. Refs. [14,152]. Field studies that reported mattress dust loads are listed in **Table 1**, which are typically in the range of  $0.1\text{--}1$   $\text{g/m}^2$ . These values are comparatively less than dust loads on indoor flooring materials, which are often between  $1$  and  $10$   $\text{g/m}^2$  [14]. Daily vacuuming of the mattress has been shown to reduce dust loading, from  $1.5$   $\text{g/m}^2$  initially, to  $0.27$   $\text{g/m}^2$  after 8 weeks of vacuuming [173].

There have been no published studies that have explicitly measured the size distribution of settled mattress dust, although it likely spans over four orders of magnitude, from sub- $1$   $\mu\text{m}$  to  $>100$   $\mu\text{m}$  particles, given the diverse spectrum of biological and inorganic particles that can exist: allergen particles, fungal spores and fragments, bacterial cells and their agglomerates, squames, bedding fiber fragments, dust mite bodies ( $\sim 200\text{--}300$   $\mu\text{m}$ ), and particles originating elsewhere in the residence that have deposited on, or tracked-in to, the bed. Mite and animal allergens, such as Der p 1, Der f 1, Fel d 1, Bal g 2, Mus m 1 (mouse), and Rat n 1 (rat), are primarily carried on particles  $\sim 5\text{--}10$   $\mu\text{m}$  in size, although their size distribution can range from  $<1$  to  $>20$   $\mu\text{m}$  [25,34,116,120]. [126] measured the size of various fungal species and found that the spores are typically in the range of  $1\text{--}4$   $\mu\text{m}$  in aerodynamic diameter. Smaller fungal fragments less than  $1$   $\mu\text{m}$ , are also common [127]. Górný et al. and Meklin et al. [58,105] reported the size distribution of various bacteria, finding aerodynamic diameters to range from  $1$  to  $>7$   $\mu\text{m}$ . It is likely many single bacterial cells exist as larger agglomerates in settled dust,  $2\text{--}4$   $\mu\text{m}$  in aerodynamic diameter, e.g. Ref. [11]. As discussed in Ref. [168]; human body sheds about  $5 \times 10^8$  skin cells per day, a significant fraction of which will accumulate in mattress dust during extended sleep periods, with the average skin flake about  $40 \times 30 \times 2$   $\mu\text{m}$  in size.

## 4. Chemical emissions from mattresses, pillows, and bed/crib frames

Although there is a robust body of literature on compositional analysis of mattress dust, there is comparatively less research on characterizing emissions of chemical contaminants from mattresses, pillows, and bed/crib frames. Several recent studies have found that infant crib mattresses are a source of a variety of chemical additives, including VOCs, plasticizers, flame retardants, and unreacted isocyanates [15,17,95], which can enter the infant sleep microenvironment through volatilization as they partition from the material-phase to the gas-phase. **Table 2** provides an

overview of chemical contaminants detected in bedding products and their associated material-phase concentrations and area-specific emission rates (SERs) to air ( $n = 14$ ).

### 4.1. Volatile organic compounds (VOCs)

The composition of a typical crib mattress includes a thick layer of polyurethane or polyester foam padding (inner-springs are also used) encased within a thin, waterproof plastic cover to protect the mattress foam and to provide an easy-to-clean surface. Studies by Refs. [3,4,15] have shown that infant crib mattresses and mattress covers are a source of a variety of VOCs, as presented in **Table 2**. Additionally, bed frames and wooden cribs may release formaldehyde, due to its presence in the resin of composite wood products used to construct the bed structure [100]. Low-VOC emitting bedding products can be found through numerous certification programs around the world, such as GREENGUARD [157], STANDARD 100 by OEKO-TEX® [115], CertiPUR-US® [24], and the Finnish M1 Emission Class for Building Materials [124].

### 4.2. Flame retardants

Crib mattresses manufactured with polyurethane foam are flammable and flame retardants are added so that they meet various flammability standards. To be effective, flame retardants are often present in the finished product at percent to tens-of-percent levels by weight [32,144,145]. Concerns over flame retardant additives in children's sleep products arose in the 1970s, when researchers demonstrated that children who wore sleepwear impregnated with the flame retardant tris(2,3-dibromopropyl) phosphate (tris-BP) were dermally exposed to the compound [13]. Throughout the next few decades, polybrominated diphenyl ether (PBDE) flame retardants, specifically pentaBDE congeners, were commonly added to crib mattress foam. Crib mattresses containing PBDEs may play a role in the elevated exposures that have been reported among infants relative to adults [61,155]. Concern over the negative health impacts of pentaBDE flame retardants led industry to end production in the U.S. in 2004. Although no longer found in new crib mattresses, these persistent compounds are still present in many older mattresses and can continue to volatilize and enter the indoor environment. This is especially important considering the long lifetime of a crib mattress ( $\sim 10$  years) and their common reuse in families with multiple children. Flame retardants currently added to crib mattresses as pentaBDE replacements include organophosphates, such as TDCPP, TCEP, and TCPP, and the Firemaster® 550 mixture [145]; **Table 2**. Recently, the State of California revised their residential flammability standard, Technical Bulletin 117 (TB-117), to require that all furniture, including mattresses for infants and adults, be sold with a label indicating the use of a chemical flame retardant additive (CA TB117-2013) [32].

### 4.3. Plasticizers

Most crib mattresses have a plastic cover for waterproofing and antibacterial purposes, and phthalates are extensively used as plasticizers to enhance the softness and flexibility of the cover. Because they are not chemically bound to the polymer matrix, they slowly volatilize from the material and migrate into surrounding environments [94,95,174,176,198]. Phthalate plasticizers identified in crib mattress covers include bis(2-ethylhexyl) phthalate (DEHP), bis(2-ethylhexyl) isophthalate (iso-DEHP), diisononyl phthalate (DINP) (**Table 2**).

Recently, the Consumer Product Safety Improvement Act was enacted in the U.S., restricting the use of certain phthalates (i.e., di-n-butyl-phthalate (DnBP), butyl-benzyl-phthalate (BBP), and

**Table 2**

Chemical contaminants detected in bedding products and their material-phase concentrations and area-specific emission rates (SERs) (n = 14).

Study	Bedding Product	Identified Chemical Contaminants in Product or Emissions to Air	Material-Phase Concentration in Product (mg/g) <sup>a</sup>	SER to Air (µg/m <sup>2</sup> h)
1. Anderson and Anderson (1999) [4]	Crib mattress covers	<b>Air:</b> toluene; 2,6-dimethyl-3-heptene; m-xylene; 1-methyl-cis-4-ethylcyclohexane; 1,3,5-trimethylbenzene; phenol; trichloroethylene; ethylbenzene; o-xylene; isopropylbenzene; diisobutylene	Not reported	Not reported
2. Anderson and Anderson (1999) [3]	Crib mattresses: foam and cover	<b>Air:</b> styrene; isopropylbenzene; limonene; ethylbenzene; 1,3-p-menthadiene; 1,3-dichlorobenzene; dipentene; 1,2,4-trimethylbenzene; nitrobenzene; 1-methyl-2-ethylbenzene; β-ocimene	Not reported	Not reported
3. Krone et al. (2003) [81]	Mattress pad foam	<b>Product:</b> total reactive isocyanate group <b>Product:</b> 2,4-toluene diisocyanate	$19 \times 10^{-3}$ $20.3 \times 10^{-3}$	Not reported
4. Kemmlein et al. (2003) [75]	Mattress Upholstery foam	<b>Air:</b> TCPP <b>Air:</b> TCPP	Not reported	23 °C: 0.012 23 °C: 77
5. Salthammer et al. (2003) [135]	Soft upholstery foam	<b>Air:</b> 1-chloro-2-propanol <b>Air:</b> 1,2-dichloropropane <b>Air:</b> TVOCs	Not reported	23 °C: 150 (24 h) 23 °C: 10 (24 h) 23 °C: < 10 (168 h)
6.-7. Hillier et al. (2003, 2009) [64,65]	Mattress foam	<b>Air (selected compounds):</b> dichloromethane; 2-methyl-2-propenal; hexane; 1-butanol; toluene; styrene; 1,3-dioxane; 1,4-dioxane; dodecane; DEP; methylcyclopentane; 3-methyl hexane; trimethylpentene; m-xylene; p-xylene; ethylbenzene; 3-methyl-nonane; limonene; ethanol; butanal; ethyl acetate; heptane; trichloroethylene; trimethyl-silanol; siloxane; tetrakis(trimethylsiloxy)silane; dodecamethyl; phenyl methyl ester formic acid; benzaldehyde; tetrachloroethene; 4-methyl-morpholine; 2,2,1,3-dioxolane; 3-methyl-pentane; 2-propenal; butanamine; n,n dimethylethanolamine; 1-chloro-2-propanol <b>Air:</b> formaldehyde	Not reported	Not reported
8. Madsen and Gibson (2008) [100]	Crib	<b>Air:</b> formaldehyde	Not reported	Emission factors (µg/h per crib): 5–3680
9. Stapleton et al. (2009) [144]	Mattress pad foam Sofa bed foam Futon foam Pillow foam	<b>Product:</b> TDCPP <b>Product:</b> TDCPP <b>Product:</b> PentaBDE <b>Product:</b> TDCPP	12 13 5 28	Not reported
10. Stapleton et al. (2011) [145]	Infant sleep positioners Crib mattresses Nursing pillows	<b>Product:</b> TCEP, TCPP, TDCPP, V6 <b>Product:</b> TCEP, TCPP, TDCPP, V6, U-OPFR, TPP, TBB/TBPH, PentaBDE <b>Product:</b> TCEP, TCPP, TDCPP, V6	>1	Not reported
11. Vangronsveld et al. (2013) [162]	Flexible polyurethane foam	<b>Product and Air:</b> 2,4-toluene diisocyanate <b>Product and Air:</b> 2,6-toluene diisocyanate	$11 \times 10^{-6}$ $45 \times 10^{-6}$	n.d.
12. Boor et al. (2014) [15]	Crib mattresses: foam and cover	<b>Air:</b> phenol  <b>Air:</b> isooctanol  <b>Air:</b> neodecanoic acid  <b>Air:</b> 2-ethyl-hexanoic acid  <b>Air:</b> 3-methyl-1-heptanol  <b>Air:</b> d-limonene  <b>Air:</b> 2,6-bis(1,1-dimethylethyl)-4-(1-oxopropyl)phenol  <b>Air:</b> (s)-3-ethyl-4-methylpentanol  <b>Air:</b> linalool  <b>Air:</b> nonanal	Not reported	23 °C: < 1–62 36 °C: 3–257 23 °C: < 1–6 36 °C: 4–7 23 °C: 3–22 36 °C: 9–40 23 °C: < 1–55 36 °C: 5–213 23 °C: 7–21 36 °C: 7–22 23 °C: 4–11 36 °C: 9–18 23 °C: 4–14 36 °C: 12–61 23 °C: 2–6 36 °C: 3–9 23 °C: 3–41 36 °C: 10–44

Table 2 (continued)

Study	Bedding Product	Identified Chemical Contaminants in Product or Emissions to Air	Material-Phase Concentration in Product (mg/g) <sup>a</sup>	SER to Air (µg/m <sup>2</sup> h)
		<b>Air:</b> decanal		23 °C: < 1–5 36 °C: 2–10
		<b>Air:</b> isopropyl myristate		23 °C: < 1–5 36 °C: 2–10
		<b>Air:</b> palmitic acid		23 °C: < 1–3 36 °C: 3–11
		<b>Air:</b> 2-ethylhexanol		23 °C: 2–10 36 °C: 12–43
		<b>Air:</b> TVOCs		23 °C: 3–6 36 °C: 7–8
13. Liang and Xu (2014b) [95]	Crib mattress covers	<b>Air:</b> DINCH	Not reported	23 °C: 87.1 (new), 22.1 (used) 36 °C: 218.8 (new), 52.1 (used)
		<b>Air:</b> DEHA		25 °C: 0.3 36 °C: 0.7
14. Boor et al. (2015b) [17]	Crib mattresses: foam and cover	<b>Product:</b> DEHP	Cover: 104.4 (new), 21 (used) Foam: 1.7 (new), 5.5 (used)	25 °C: 0.4 36 °C: 4
		<b>Product:</b> iso-DEHP	Cover: 314.5 (new), 60 (used) Foam: n.d. (new), 14.4 (used)	Not reported
		<b>Product:</b> DINP	Cover: 28.5 (new), 136.8 (used) Foam: n.d. (new), 22.3 (used)	
		<b>Product:</b> DINCH	Cover: 53.2 (new), 103.1 (used) Foam: 0.1 (new), 0.1 (used)	
		<b>Product:</b> DEHA	Cover: 4.8 (new), 31.1 (used) Foam: n.d. (new), 0.2 (used)	
		<b>Product:</b> PentaBDE	Foam: 7.46 (used)	
		<b>Product:</b> NCO	Not reported	

Nomenclature: **n.d.**: not detected; **TVOCs**: total volatile organic compounds; **DEHP**: Bis(2-ethylhexyl) phthalate; **DEP**: diethyl phthalate; **iso-DEHP**: Bis(2-ethylhexyl) isophthalate; **DINP**: Diisononyl phthalate; **DINCH**: Diisononyl 1,2-cyclohexanedicarboxylic acid; **DEHA**: Bis(2-ethylhexyl) adipate; **TCEP**: tris-(2-chloroethyl)-phosphate; **TCPP**: tris-(1-chloro-2-propyl) phosphate; **TDCPP**: tris-(2,3-dichloropropyl)-phosphate; **TPP**: triphenyl phosphate; **V6**: 2,2-bis(chloromethyl)propane-1,3-diyl-tetrakis(2-chloroethyl)bis(phosphate); **PentaBDE**: pentabromodiphenyl ether; **U-OPFR**: 2,2-bis(chloromethyl)propane-1,3-diyl tetrakis(1-chloropropan-2-yl)bis(phosphate); **TBB**: 2-ethylhexyl-2,3,4,5-tetrabromobenzoate; **TBPH**: bis(2-ethylhexyl)-3,4,5,6-tetrabromophthalate; **NCO**: unreacted isocyanates.

<sup>a</sup> mg of compound per g of material.




Fig. 1. Duration of time (h/day) spent in sleep or nap activity [159]; data ID = 14500). Box plots represent interquartile range and whiskers represent the 5th and 95th percentiles.

DEHP) in toys and child care articles (66). However, there is some debate as to whether or not crib mattresses are included in the definition of “child care articles” [158,201]. In addition, the ban on DINP and di-iso-decyl-phthalate (DIDP) only applies to children's toys that can be placed in the mouth. Alternative plasticizers that have a chemical structure similar to phthalates, such as di(2-ethylhexyl) adipate (DEHA) and diisononyl cyclohexane-1,2-dicarboxylate (DINCH), have emerged recently and have been identified in new crib mattress covers [17]; Table 2.

#### 4.4. Unreacted isocyanates

An excess of toluene diisocyanate (TDI, the predominate diisocyanate used in polyurethane foam production [162]), may be added during the manufacture of polyurethane foam, above which

is necessary for it to react with hydroxyl groups of polyols and water [185]. As such, unreacted isocyanate may be present in the final polyurethane foam product [20,81,185]. Krone et al. [81] identified NCO in polyurethane foam-containing consumer products, including a mattress and sofa padding. NCO was also detected in polyurethane-foam containing crib mattresses via PAS-FT-IR analysis in Ref. [17]. The foam in crib mattresses is somewhat protected from excess moisture released from the infant body, which can react with the isocyanates [10].

#### 5. The source-proximity effect of the sleep microenvironment

In the sleep microenvironment, there are several key factors that contribute to the source-proximity effect (Fig. 5), including the spatial proximity of the BZ to the source, incomplete mixing and poor ventilation of room air, concentrations gradients near an actively emitting source, the personal cloud due to human body movement-induced particle resuspension, the buoyant human thermal plume, heat transfer from the human body to the source, which may elevate the emissions of gaseous pollutants, and direct dermal contact with the source.

##### 5.1. Spatial proximity and concentration gradients near actively emitting sources

The spatial proximity of a person's BZ to their mattress and bedding can lead to elevated concentrations in their BZ relative to the bulk bedroom air. The spatial proximity effect has been evaluated in numerous studies, which typically release inert tracer gases from point or area sources and use sampling arrays to measure the uniformity of the dispersion of the tracer gas in a space, e.g. Refs. [1,83,103]. Collectively, these studies have shown that tracer gas concentrations near actively emitting sources are greater than those at locations further from the source, typically by a factor greater than 2.




Fig. 2. Fraction of time adults spent sleeping or napping in a residence [77,159].




Fig. 3. Volume of air inhaled during sleep or nap activity per day, normalized by body mass for each age group and gender (calculated using U.S. EPA EFH data set, [159]).

Laverge et al. [88] highlighted the important role of the spatial proximity effect on inhalation exposure to pollutants originating in the sleep microenvironment. A breathing thermal manikin was positioned on a twin-size mattress with a pillow, from which sulfur hexafluoride (SF<sub>6</sub>) was uniformly released to simulate the emission of gaseous pollutants. The relative intake fraction, defined as the ratio of the BZ concentration to the well-mixed chamber concentration, was reported for different sleeping positions and bedding arrangements. BZ SF<sub>6</sub> concentrations were found to be significantly greater than those measured in the bulk air, with mean relative

intake fractions for the mattress source of 1.24 for the supine position, 1.39 for the lateral position, and 1.73 for the prone position. Higher mean relative intake fractions were reported for the pillow source with 2.16 for the supine position and between 3.39 and 4.02 for the lateral position. Additionally, covering the entire manikin's body and head with a blanket increased the relative intake fraction to 32.7. Boor et al. [15] reported similar findings for VOCs emitted from a crib mattress, with BZ TVOC concentrations greater than those in the bulk chamber air by factors in the range of 1.8–2.4. In regard to resuspended mattress dust particles, [16] also found BZ




Fig. 4. Infant dermal exposure dose considerations in the sleep microenvironment: body surface area normalized by body mass (BSA/BW, m<sup>2</sup>/kg) for each age group and gender (calculated using U.S. EPA EFH data set, [159]).

concentrations to be greater than bulk air concentrations by factors of 1.07–1.94, depending on particle size. The studies demonstrate that a person will receive elevated exposures to particulate and gaseous pollutants that are released from their mattresses, beyond what would be estimated from bulk bedroom air measurements.

### 5.2. Incomplete mixing of bedroom air

The spatial proximity effect is due in part to incomplete mixing of the released pollutant with the bulk air, which leads to the development of concentration gradients in the space, as suggested in some of the aforementioned studies. Additional studies by Refs. [18,41,49,101,129,166,178]; among others, have found that incomplete mixing conditions are associated with non-uniform pollutant concentration distributions in indoor environments.

Bedrooms, when compared to the rest of the home, have been found to be poorly ventilated, e.g. Refs. [6,8,90,199]. Incomplete mixing may be aided by the low ventilation rates that have been reported for bedrooms and the reduced run time of ventilation systems at night. Bekö et al. [8] measured the ventilation rates in the naturally ventilated bedrooms of 500 children in Denmark and reported a mean ventilation rate of  $0.46 \text{ h}^{-1}$ , with rates as low as  $0.1 \text{ h}^{-1}$ . In addition, [147] found that the fractional operation time of residential ventilation systems is the lowest during the late evening/early morning hours (10PM to 7AM). As the ventilation system run time is reduced, a residence relying on forced mechanical ventilation to deliver fresh (or filtered) air to the bedroom during sleep periods may see poor mixing conditions in the space (e.g. more stagnant air due to reduced ventilation and less human activity and movement to enhance mixing) and inadequate dilution of pollutants released within the sleep microenvironment, leading to the accumulation of pollutants in the air. Furthermore, it is expected that for an actively emitting diffusion-controlled source, such as a mattress, concentrations in very close proximity (several cm) will be greater than those in the bulk air, as a chemical compound partitions from the source to the air immediately above it. Thus, it cannot be assumed that a pollutant released from a mattress or bedding will be well mixed with the bulk bedroom air, and BZ concentrations should be measured to accurately estimate inhalation exposure to pollutants originating in this microenvironment.

### 5.3. Personal cloud effect and human-induced resuspension of mattress dust

The personal cloud effect, whereby people can be envisioned as being surrounded by a cloud of particles, many of which are of

biological origin, is another factor that may lead to elevated exposures in the sleep microenvironment [12,96,104]. The personal cloud effect was used to explain the consistently higher personal exposures to particles compared to stationary monitoring locations [165]. Rabinovitch et al. [122] provides one example in which personal (approximate BZ) bacterial endotoxin concentrations measured with personal monitors are about 3 times greater than those measured at stationary indoor locations, suggesting that endotoxin exposures are better correlated with sources in close proximity to the person and their BZ than those in the bulk indoor air. Human-induced particle resuspension from indoor surfaces, including mattresses and bedding, is one of the physical processes contributing to the personal cloud effect, leading to elevations in BZ concentrations beyond those in the bulk air. In the sleep microenvironment, the personal cloud effect is likely to be enhanced due to the close proximity of a person's BZ to the surfaces from which particle resuspension is occurring, an important consideration for larger particles ( $>10 \mu\text{m}$ ) that have high settling velocities (on the order of  $\sim 10^{-3} \text{ m/s}$ ), and likely short trajectories upward from a surface. Investigations by Spilak et al. and Boor et al. [16,141] have shown that human body movements in bed, such as rolling from the prone to supine position, can resuspend significant quantities of settled particles on mattresses, pillows, bed sheets, and blankets. Spilak et al. [141] demonstrated that a pillow cover (225-thread count) does not serve as an effective barrier to the transport of resuspended particles from the underlying pillow surface, however, the impact of impermeable mattress covers on particle resuspension and transport to the BZ has yet to be investigated.

### 5.4. Body movements during sleep periods

Mattress dust resuspension is driven by human body movements in bed. Human activity in bed can occur during periods of wakefulness, e.g. as a person enters a bed and prepares themselves for sleep, and throughout their extended sleep state. Although the absence of voluntary motor behavior is a characteristic of the sleep state, movements are commonly reported, ranging from about ten significant body posture shifts (e.g. Ref. [186]), to several hundred smaller body movements (e.g. Ref. [187]). The cause of body movements during sleep periods is largely unknown, although several theories suggest that movements occur to relieve muscular discomfort during extended periods of immobility [67,188].

Sleep is typically divided into several sleep stages: stage W (Wakefulness), NREM stage 1 (non-rapid eye movement), NREM stage 2, NREM stage 3 or SWS (Slow Wave Sleep, e.g. "deep sleep"), and REM (rapid eye movement) (e.g. Ref. [189]). It is important to understand the dependency of body movements on sleep stages, as




Fig. 5. The source-proximity effect of the sleep microenvironment.

body movements are responsible for inducing the necessary removal forces (aerodynamic, mechanical) to resuspend mattress dust particles [121]. Shimohira et al. [190] found a strong dependency of body movement intensity on sleep stage among children 4–12 years of age. Body movements were classified into four groups, GM-1: axial rotation of the body, GM-2: movement of limbs and trunk (torso) muscles without rotation, GM-3: movement of two or more limbs but not trunk muscles, and GM-4: movement of only one limb. The authors found that movements are much more frequent during both NREM stage 1 and REM sleep, with a decrease during SWS (deep sleep). The most commonly reported movement was GM-1, in which the child rotates some degree, with up to 13 movements/hour during NREM 1 sleep.

Wilde-Frenz and Schulz [191] observed a similar trend among adults, 21–36 years of age, where body movement frequency was the greatest during stage W, followed by NREM 1, then REM, then NREM 2, and SWS. Body movement frequency ranged from <0.1 to 0.8 body movements per sleep minute (bm/min), depending on the sleep stage. Giganti et al. [188] found that the number of body movements were greatest during NREM 1 sleep, with movement frequency remaining nearly constant during NREM 2, SWS, and REM sleep. Body movements were found to decrease with age, with an average of 0.25 bm/min for age group 1 and 0.09 bm/min for age group 2, equating to an average of 86 body movement during the entire sleep period for group 1 (mean sleep time of 342 min) and 27 movements for group 2 (mean sleep time of 297 min).

The findings of these studies suggest that body movement frequency is a function of both sleep stage and age. Thus, the potential for human-induced particle resuspension would vary throughout a sleep period, and may be greater during periods of more frequent activity (e.g. NREM 1 and REM stages) than during deep sleep stages (SWS). [16] demonstrated that more intense body movements in bed, such as a full 360° rotation of the torso, are associated with higher surface vibrations and emissions of resuspended particles compared to less intense movements (e.g. sitting on the mattress). Studies have suggested that asthmatic children and adults have more disturbed sleep patterns [192,193], and Zanobetti et al. [194] found sleep-disordered breathing, defined as the recurrent episodic disruption of normal breathing during sleep, and decreases in sleep efficiency to be associated with increases in short-term variation in the airborne concentration of 10 µm particles. People with asthma and other breathing ailments may have a greater frequency of body movements during their disrupted sleep period, which may put them at greater risk for elevated exposure to resuspended mattress dust particles.

### 5.5. Human thermal plume

In the sleep microenvironment, the airflow pattern is a result of two different flows and their interaction: the buoyant human thermal plume and transient flows generated by breathing [195,196]. While the thermal plume depends on the metabolic rate of a sleeping person and the sleep stage, the transient flow is mainly dependent on whether the sleeping person inhales/exhales via nose or mouth. The buoyant human thermal plume may also be responsible for elevated BZ concentrations for pollutant sources in close proximity of the human body. Rim and Novoselac [128] and Rim and Novoselac [129] demonstrated that the thermal plume is an effective mechanism for transporting pollutants, released from various locations in close vicinity to the human body, vertically upward, toward to BZ. Thus, the thermal plume that develops around a sleeping person may aid in transporting pollutants released from the mattress surface, upward and toward their BZ. However, the thermal plume may also serve the secondary role of entraining “fresh” air exterior to the sleep microenvironment,

thereby diluting concentrations of mattress pollutants in the BZ, as suggested by Ref. [88]. The thermal plume also interacts with the shape of the breathing zone, especially the inhalation zone, reducing it to a shallow area of only a few centimeters wide stretching down around the cheeks of a person resting in supine position [89].

### 5.6. Impact of body temperature and sweating on chemical emissions from mattresses

Another unique attribute of the sleep microenvironment that may play an important role in enhancing the source-proximity effect is the warming of nearby objects via heat exchange with the human body. Along with warming the surrounding air via convective heat exchange, leading to the development of the thermal plume, our bodies will also transfer heat to surfaces that are in direct contact via conduction and those in close proximity via radiation. Lu et al. [99] monitored the surface temperature of a mattress around sleeping adult male subjects with a matrix of 16 thermistors. Maximum surface temperatures were found to range from 31 to 36 °C beneath and around the lower limbs and between 34 and 36 °C beneath and around the waist. Thus, it is to be expected that the sleeping human body will increase the surface temperature of their mattress, pillow, and bedding materials. The emissions of gaseous pollutants, such as VOCs and SVOCs, from building materials has been shown to increase with temperature (e.g. Refs. [98,197,198]), with recent studies by Refs. [15,95] observing this effect in crib mattresses and mattress covers. Thus, the close proximity of the human body to potential pollutant sources may enhance volatilization, and subsequently, further increase BZ concentrations. Throughout the sleep period, the sleeping thermal environment is a permanently humid environment due to the production of large quantities of moisture through sweating and the large buffering capacity provided by mattresses and bedding materials. Elevated moisture levels may also have a strong impact on emissions of VOCs and SVOCs from mattresses, as well as their desorption from bedding materials that can serve as sorptive reservoirs during the day (e.g. Ref. [112]).

### 5.7. Direct dermal contact with mattresses and bedding materials

Transdermal uptake of pollutants, such as SVOCs, via contact transfer and air-to-skin transport, is another contributing factor to the source-proximity effect. During sleep periods, a person's body, and exposed skin, is in intimate contact with their pajamas, mattress, mattress cover, and bedding materials. Dermal exposure is an important pathway for certain SVOCs, including plasticizers that are added to mattress covers [109,169]. If the gas-phase concentrations of these pollutants are greater near the source or the skin, dermal uptake via air-to-skin transport may be enhanced [108]. Laverge et al. [88] demonstrated that concentrations of mattress-released gaseous pollutants (SF<sub>6</sub>) under the bedding (where most of the body is exposed) can be up to 30 times higher than those in the bulk air of the room. Additionally, during the day, SVOCs found in mattresses, as well as in the bulk bedroom air, may partition to bedding fabric fibers and pajamas [108]. Fabric fibers may also be coated with a surface film of detergents and fabric softeners that are used for cleaning. Temperature elevations during sleep periods may promote desorption of the sorbed SVOCs, thereby offering an additional route of transdermal uptake [169].

### 5.8. Personalized ventilation and air cleaning systems for beds

An important factor that can affect pollutant transport dynamics and exposures while sleeping is the use of personalized ventilation

(PV) and air cleaning systems for beds. Fan-forced decentralized ventilation systems commonly found in homes can provide filtered and conditioned air for a sleeping person, e.g. Ref. [90]. However, they may not necessarily be sufficient in reducing concentrations of mattress-released pollutants within the sleep microenvironment or in improving sleep quality among people with asthma and allergies [56,59,76]. PV systems can deliver a relatively small amount of air (less than  $0.3 \text{ h}^{-1}$ ) directly to the BZ of a sleeping person, thereby offering an alternative strategy to reduce sleep exposures. Several PV designs for beds have been introduced, including: bedside PV [84]; overhead PV [106]; and a temperature-controlled laminar airflow (TLA) PV [143]. Melikov et al. [106] found overhead PV to reduce exposures of medical staff to the exhaled air of a patient in a bed at significantly lower ventilation rates compared to a decentralized ventilation system. Spilak et al. [143] found TLA PV to reduce BZ concentrations of airborne particles with a sleeping thermal manikin beyond what can be achieved with a portable air cleaner. Furthermore, TLA PV has been shown to reduce the severity of asthma symptoms among patients with persistent asthma [19,118,136]. Along with PV, portable HEPA filtration devices positioned close to the bed (e.g. on a night-stand) have been shown to reduce BZ concentrations of coarse and fine airborne particles [28,142].

## 6. Conclusions and future directions

We spend nearly one-third of every day asleep. Indoor air quality (IAQ) in the sleep microenvironment, and more broadly, in bedrooms, may have important implications for human health and well-being, especially among children and adults with asthma and allergies. Human exposures to mattress-released pollutants can be augmented as air is inhaled just centimeters above the source, with breathing zone concentrations typically greater than those in the bulk air by a factor of two, and in specific conditions (cover over head), as high as 30. Beds serve as reservoirs for a complex mixture of biological- and chemical-laden dust, which can be readily stirred-up upon tossing and turning during sleep. House dust mite allergens and fungi may be generated in-situ, while other particles, such as bacterial cells and animal allergens, may be generated by, or tracked-in with, people and their pets. The sleep thermal environment can promote fungal growth and the proliferation of dust mites. Settled dust particles can accumulate over time, especially given the low cleaning frequencies of mattresses and pillows. Exposure to SVOCs via inhalation and dermal pathways during sleep may be significant, given the ubiquity of SVOC additives in infant and adult bedding products (foams and plastic covers) and their enhanced volatilization in and under bedding due to elevated localized surface temperatures ( $\sim 30 \text{ }^\circ\text{C}$ ) and relative humidities.

This review provided an overview of literature pertaining to the emerging field of what can be referred to as *sleep IAQ*, the study of indoor air pollutant dynamics and exposures in sleep microenvironments. However, the existing research on *sleep IAQ* is insufficient to elucidate the impact of sleep exposures to indoor air pollutants and biological material on human health and sleep quality. In reviewing the extant literature, the following research needs were identified:

- (1.) Field studies are needed to understand the variation in exposures occurring during sleep among diverse populations and age-groups and how these exposures compare to exposures occurring in the remaining two-thirds of the day. In particular, the contribution of early-life sleep exposures to microbes, allergens, and SVOCs to total exposures to these agents should be evaluated. This work would necessitate personal exposure monitoring (BZ sampling) of particles and

gases. Improved sleep exposure assessment will permit more reliable associations with health outcomes, beyond what can be achieved through basic mattress dust sampling and analysis.

- (2.) Research is needed to elucidate the impact of human exposure to mattress-released pollutants, human bioeffluents, and pollutants originating elsewhere in the residence or outdoors on sleep itself. Several recent studies have indicated that poorly ventilated bedrooms can be associated with higher  $\text{CO}_2$  concentrations, which can adversely affect sleep quality and next day performance [87,148], and that exposure to traffic pollution (e.g. black carbon) may disturb sleep (e.g. Refs. [48,55]). These efforts can be extended to address the full range of particulate and gaseous indoor air pollutants.
- (3.) Little is known on the role of multiple bed occupants and pets in influencing sleep exposures and pollutant transport dynamics. According to the National Sleep Foundation (NSF) Sleep in America Poll, 48–72% of Americans sleep with their significant other and 2–16% sleep with their pets, depending on ethnicity [111]. Multiple people or pets in bed may impact the microbial composition of mattress dust and the extent to which it is aerosolized throughout the night, as well as the interaction of multiple buoyant thermal plumes in the air in and around the bed.
- (4.) Research in the *sleep IAQ* domain should follow a mechanistic approach and include new research on the following topics: experimental and modeling studies on the emissions of VOCs and SVOCs from new and used bedding products on the market, including eco-friendly mattresses and low-cost air mattresses; transient sorption-desorption cycling of these chemical contaminants with bedding and pajama fabrics during periods of occupancy and non-occupancy in bed, as the thermal environment changes with time; size-resolved resuspension fractions of biological particles from new and soiled bedding materials; and improved assessment of inter-zonal airflow between the BZ and bulk bedroom air under a range of bedroom ventilation conditions.
- (5.) Future investigations should explore exposures in unique sleep microenvironments, such as neonatal intensive care units (NICUs), hospital beds, tents for camping (which can contain high levels of flame retardants, [74]), hotel beds, and bedrooms with many occupants, such as shared dormitory rooms.

## Acknowledgements

Financial support was provided by Purdue University start-up funds, the U.S. Environmental Protection Agency Science to Achieve Results Fellowship Program (Grant No. F13D10740), and the National Science Foundation (Grant No. CBET-1150713).

## References

- [1] V. Acevedo-Bolton, K.-C. Cheng, R.-T. Jiang, W.R. Ott, N.E. Klepeis, L.M. Hildemann, Measurement of the proximity effect for indoor air pollutant sources in two homes, *J. Environ. Monit.* 14 (2012) 94–104.
- [2] N. Ali, A.C. Dirtu, N. Van den Eede, E. Goosey, S. Harrad, H. Neels, A. t Man-netje, J. Coakley, J. Douwes, A. Covaci, Occurrence of alternative flame retardants in indoor dust from New Zealand: indoor sources and human exposure assessment, *Chemosphere* 88 (2012) 1276–1282.
- [3] R.C. Anderson, J.H. Anderson, Respiratory toxicity of mattress emissions in mice, *Arch. Environ. Health* 54 (1999) 202–209.
- [4] R.C. Anderson, J.H. Anderson, Respiratory toxicity in mice exposed to mattress covers, *Arch. Environ. Health* 55 (2000) 38–43.
- [5] C. Barnig, G. Reboux, S. Rousset, A. Casset, C. Sohy, J.-C. Dalphin, F.de Blay, Indoor dust and air concentrations of endotoxin in urban and rural environments, *Lett. Appl. Microbiol.* 56 (2012) 161–167.



- [6] P. Batog, M. Badura, Dynamic of changes in carbon dioxide concentration in bedrooms, in: The 11th International Conference on modern building materials, structures and Techniques, MBMST 2013, 2013.
- [7] M. Begum, A. Aya, C. Cheong, H. Neumeister-Kemp, K. White, P. Kemp, Prevalence of Fungi in Used Pillows, in: Proceedings of Healthy Buildings 2012, 2012. Brisbane, Australia, paper ID: 5H.4.
- [8] G. Bekö, T. Lund, F. Nors, J. Toftum, G. Clausen, Ventilation rates in the bedrooms of 500 Danish children, *Build. Environ.* 45 (2010) 2289–2295.
- [9] G. Bekö, G. Morrison, C.J. Weschler, H.M. Koch, C. Pälmeke, T. Salthammer, T. Schripp, J. Toftum, G. Clausen, Measurements of dermal uptake of nico ne directly from air and clothing, *Indoor Air* 27 (2017) 427–433.
- [10] D. Bello, C.A. Herrick, T.J. Smith, S.R. Woskie, R.P. Streicher, M.R. Cullen, Y. Liu, C.A. Redlich, Skin exposure to isocyanates: reason for concern, *Environ. Health Persp* 115 (2007) 328–335.
- [11] S. Bhangar, J.A. Huffman, W.W. Nazaroff, Size-resolved fluorescent biological aerosol particle concentrations and occupant emissions in a university classroom, *Indoor Air* 24 (6) (2014) 604–617.
- [12] S. Bhangar, R.I. Adams, W. Pasut, J.A. Huffman, E.A. Arens, J.W. Taylor, T.D. Bruns, W.W. Nazaroff, Chamber bioaerosol study: human emissions of size-resolved fluorescent biological aerosol particles, *Indoor Air* 26 (2) (2015) 193–206.
- [13] A. Blum, M.D. Gold, B.N. Ames, C. Kenyon, F.R. Jones, E.A. Hett, R.C. Dougherty, E.C. Horning, I. Dzidic, D.I. Caroll, R.N. Stillwell, J.P. Thenol, Children absorb Tris-BP flame retardant from sleepwear: urine contains the mutagenic metabolite, 2,3-dibromopropanol, *Science* 201 (1978) 1020–1023.
- [14] B.E. Boor, J.A. Siegel, A. Novoselac, Monolayer and multilayer particle deposits on hard surfaces: literature review and implications for particle resuspension in the indoor environment, *Aerosol Sci. Technol.* 47 (8) (2013) 831–847.
- [15] B.E. Boor, H. Järnström, A. Novoselac, Y. Xu, Infant exposure to emissions of volatile organic compounds from crib mattresses, *Environ. Sci. Technol.* 48 (6) (2014) 3541–3549.
- [16] B.E. Boor, M.P. Spilak, R.L. Corsi, A. Novoselac, Characterizing particle resuspension from mattresses: chamber study, *Indoor Air* 25 (4) (2015a) 441–456.
- [17] B.E. Boor, Y. Liang, N.E. Crain, H. Järnström, A. Novoselac, Y. Xu, Identification of phthalate and alternative plasticizers, flame retardants, and unreacted isocyanates in infant crib mattress covers and foam, *Environ. Sci. Technol. Lett.* 2 (4) (2015b) 89–94.
- [18] S.M. Bowes, E.G. Mason, M. Corn, Confined space ventilation: tracer gas analysis of mixing characteristics, *AIHA J.* 54 (1993) 639–646.
- [19] R.J. Boyle, C. Pedroletti, M. Wickman, M. Bjermer, E. Valovirta, R. Dahl, A. Von Berg, O. Zetterström, J.O. Warner, Nocturnal temperature controlled laminar airflow for treating atopic asthma: a randomized controlled trial, *Thorax* 67 (2012) 215–221.
- [20] California Office of Environmental Health Hazard Assessment (CA OEHHA), Proposed Revised Reference Exposure Levels for Toluene Diisocyanate and Methylene Diphenyl Diisocyanate, CA OEHHA, Sacramento, CA, 2010.
- [21] M. Callesen, G. Bekö, C.J. Weschler, T. Sigsgaard, T.K. Jensen, G. Clausen, J. Toftum, L.A. Norberg, A. Høst, Associations between selected allergens, phthalates, nicotine, PAHs and bedroom ventilation and clinically confirmed asthma, rhinoconjunctivitis and atopic dermatitis in preschool children, *Indoor Air* 24 (2) (2014) 136–147.
- [22] D. Canbaz, M.J.M. van Velzen, E. Hallner, A.H. Zwinderman, M. Wickman, P.E.G. Leonards, R. van Ree, L.S. van Rijt, Exposure to organophosphate and polybrominated diphenyl ether flame retardants via indoor dust and childhood asthma, *Indoor Air* 26 (2016) 403–413.
- [23] L. Casas, C. Tischer, I.M. Wouters, M. Valkonen, U. Gehring, G. Doekes, M. Torrent, J. Pekkanen, R. Garcia-Esteban, A. Hyvärinen, J. Heinrich, J. Sunyer, Endotoxin, extracellular polysaccharides, and  $\beta(1-3)$ -glucan concentrations in dust and their determinants in four European birth cohorts: results from the HITEA project, *Indoor Air* 23 (2013) 208–216.
- [24] CertiPUR-US, Physical Performance and Environmental Guidelines for Certification of Slabstock Flexible Polyurethane Foam for Use in Furniture and Bedding, Edition 07.31.2017, 2017.
- [25] C. Chang, M.E. Gershwin, Indoor air quality and human health, *Clin. Rev. Allerg. Immu* 27 (2004) 219–239.
- [26] C.-M. Chen, A. Mielck, B. Fahlbusch, W. Bischof, O. Herbarth, M. Borte, H.-E. Wichmann, J. Heinrich, Social factors, allergen, endotoxin, and dust mass in mattress, *Indoor Air* 17 (2007) 384–393. *Clin. Exp. Dermatol.* 35:238–244.
- [27] C.-M. Chen, S. Sausenthaler, W. Bischof, O. Herbarth, M. Borte, H. Behrendt, U. Krämer, H.C. Williams, H.-E. Wichmann, J. Heinrich, Perinatal exposure to endotoxin and the development of eczema during the first 6 years of life, *Clin. Exp. Dermatol* 35 (2009) 238–244.
- [28] J.D. Clark, A. Novoselac, R.L. Corsi, Assessment of Two Control Strategies for Reducing Human Exposure to Airborne Particles while Sleeping in Low-Energy Homes, in: Proceedings of ASHRAE IAQ 2013, 2013. Vancouver, British Columbia, Canada.
- [29] C.A. Clayton, L.A. Wallace, H. Ozkaynak, J.D. Spengler, Particle total exposure assessment methodology (PTEAM) study: distributions of aerosol and elemental concentrations in personal, indoor, and outdoor air samples in a southern California community, *J. Expos. Anal. Envir. Epidemi* 6 (1993) 227–250.
- [30] E.A. Cohen Hubal, L.S. Sheldon, J.M. Burke, T.R. McCurdy, M.R. Berry, M.L. Rigas, V.G. Zartarian, N.C.G. Freeman, Children's exposure assessment: a review of factors influencing children's exposure, and the data available to characterize and assess that exposure, *Environ. Health Persp* 108 (6) (2000) 475–486.
- [31] W.B. Cooke, M.J. Foter, Fungi in used bedding materials, *Appl. Microbiol.* 6 (1958) 169–173.
- [32] E.M. Cooper, G. Kroeger, K. Davis, C.R. Clark, P. Lee Ferguson, H.M. Stapleton, Results from screening polyurethane foam based consumer products for flame retardant chemicals: assessing impacts on the change in the furniture flammability standards, *Environ. Sci. Technol.* 50 (2016) 10653–10660.
- [33] N.F. Conant, H.C. Wagner, F.M. Rackemann, Fungi found in pillows, mattresses and furniture, *Allergy* 7 (1936) 234–237.
- [34] A. Custovic, B. Simpson, A. Simpson, C. Hallam, M. Craven, A. Woodcock, Relationship between mite, cat, and dog allergens in reservoir dust and ambient air, *Allergy* 54 (1999) 612–616.
- [35] D.R. da Silva, R.S. Binotti, C.M. da Silva, C.H. de Oliveira, A. Condino-Neto, E.M. de Capitani, Mites in dust samples from mattress surfaces from single beds or cribs in the south Brazilian city of Londrina, *Pediatr. Allergy Immunol.* 16 (2005) 132–136.
- [36] K.C. Dannemiller, C.J. Weschler, J. Peccia, Fungal and bacterial growth indoor dust at elevated relative humidity levels, *Indoor Air* (2016), <http://dx.doi.org/10.1111/ina.12313>.
- [37] U. Diez, T. Kroebner, M. Rehwagen, M. Richter, H. Wetzig, R. Schulz, M. Borte, G. Metzner, P. Krumbiegel, O. Herbarth, Effects of indoor painting and smoking on airway symptoms in atopy risk children in the first year of life results of the LARS-study. Leipzig Allergy High-Risk Children Study, *Int. J. Hyg. Environ. Health* 203 (2000) 23–28.
- [38] J. Douwes, G. Doekes, J. Heinrich, A. Koch, W. Bischof, B. Brunekreef, Endotoxin and  $\beta(1-3)$ -glucan in house dust and the relation with home characteristics: a pilot study in 25 German houses, *Indoor Air* 8 (1998) 255–263.
- [39] J. Douwes, A. Zuidhof, G. Doekes, S.C. van der Zee, I. Wouters, M.H. Boezen, B. Brunekreef,  $(1-3)$ - $\beta$ -D-Glucan and endotoxin in house dust and peak flow variability in children, *Am. J. Respir. Crit. Care Med.* 162 (2000) 1348–1354.
- [40] V. Doyen, A.-B. Johansson, L. Hanssens, N. Dehennin, D.H.P. Dinh, G. Casimir, O. Michel, Relationship between the presence of newborn and the house dust endotoxin, *Sci. Total Environ.* 409 (2011) 5313–5317.
- [41] A.C. Drescher, C. Lobascio, A.J. Gadgil, W.W. Nazaroff, Mixing of a point-source indoor pollutant by forced convection, *Indoor Air* 5 (1995) 204–214.
- [42] B. Eberlein, A.F. Gulyas, K. Schultz, J. Lecheler, S. Flögel, C. Wolfmeyer, K. Thiessen, T. Jakob, T. Schuster, R. Hollweck, J. Ring, H. Behrendt, Domestic allergens and endotoxin in three hospitals offering in-patient rehabilitation for allergic diseases in the alpine mountain climate of Bavaria – the AURA study, *Int. J. Hyg. Environ. Health* 212 (2009) 21–26.
- [43] R.D. Edwards, J. Jurvelin, K. Saarela, M. Jantunen, VOC concentrations measured in personal samples and residential indoor, outdoor and workplace microenvironments in EXPOLIS-Helsinki, *Finl. Atmos. Environ.* 35 (2001) 4531–4543.
- [44] N. El Sharif, J. Douwes, P.H.M. Hoet, G. Doekes, B. Nemert, Concentrations of domestic mite and pet allergens and endotoxin in Palestine, *Allergy* 59 (2004) 623–631.
- [45] M.J. Ege, M. Mayer, K. Schwaiger, J. Mattes, G. Pershagen, M. van Hage, A. Scheynius, J. Bauer, E. von Mutius, Environmental bacteria and childhood asthma, *Allergy* 67 (2012) 1565–1571.
- [46] Eurostat, Time Spent, Participation Time and Participation Rate in the Main Activity by Sex and Age Group – Collection Round 2000, 2000. <http://ec.europa.eu/eurostat/en/data/database>.
- [47] B. Fahlbusch, D. Hornung, J. Heinrich, H.-M. Dahse, L. Jäger, Quantification of group 5 grass pollen allergens in house dust, *Clin. Exp. Allergy* 30 (2000) 1645–1652.
- [48] S.C. Fang, J. Schwartz, M. Yang, H.K. Yaggi, D.L. Bliwise, A.B. Araujo, Traffic-related air pollution and sleep in the Boston area community health survey, *J. Expo. Sci. Environ. Epidemiol.* 25 (5) (2015) 451–456.
- [49] E.J. Furtaw, M.D. Pandian, D.R. Nelson, J.V. Behar, Modeling indoor air concentrations near emission sources in imperfectly mixed rooms, *J. Air Waste Manag. Assoc.* 46 (1996) 861–868.
- [50] L. Gagliardi, F. Rusconi, Respiratory rate and body mass in the first three years of life, *Arc. Dis. Child.* 76 (1997) 151–154.
- [51] M.H. Garrett, B.M. Hooper, M.A. Hooper, Indoor environmental factors associated with house-dust-mite allergen (Der p 1) levels in south-eastern Australian houses, *Allergy* 53 (1998) 1060–1065.
- [52] U. Gehring, W. Bischof, M. Borte, O. Herbarth, H.-E. Wichmann, J. Heinrich, Levels and predictors of endotoxin in mattress dust samples from East and West German homes, *Indoor Air* 14 (2004) 284–292.
- [53] U. Gehring, J.C. de Jongste, M. Kerkhof, M.A. Oldewening, D. Postma, R.T. van Strien, A.H. Wijga, S.M. Willers, A. Wolse, J. Gerritsen, H.A. Smit, B. Brunekreef, The 8-year follow-up of the PIAMA intervention study assessing the effect of mite-impermeable mattress covers, *Allergy* 67 (2012) 248–256.
- [54] M.E. Giovannangelo, U. Gehring, E. Nordling, M. Oldenwening, K. van Rijswijk, S. de Wind, G. Hoek, J. Heinrich, T. Bellander, B. Brunekreef, Levels and determinants of  $\beta(1-3)$ -glucans and fungal extracellular polysaccharides in house dust of (pre-)school children in three European countries, *Environ. Int.* 33 (2007) 9–16.
- [55] T. Gislason, R.J. Bertelsen, F.G. Real, T. Sigsgaard, K.A. Franklin, E. Lindberg,

- C. Janson, E.S. Arnardottir, J. Hellgren, B. Benediktsdottir, B. Forsberg, A. Johannessen, Self-reported exposure to traffic pollution in relation to daytime sleepiness and habitual snoring: a questionnaire study in seven North-European cities, *Sleep, Med.* 24 (2016) 93–99.
- [56] P.C. Goetzsche, H.K. Johansen, House dust mite control measures for asthma, *Cochrane Database Syst. Rev.* 18 (4) (2008) CD001187.
- [57] R.B. Gore, E.A. Hadi, M. Craven, F.I. Smillie, T.J. O'Meara, E. Tovey, A. Woodcock, A. Custovic, Personal exposure to house dust mite allergen in bed: nasal air sampling and reservoir allergen levels, *Clin. Exp. Allergy* 32 (2002) 856–859.
- [58] R.L. Górny, J. Dutkiewicz, E. Krysińska-Traczyk, Size distribution of bacterial and fungal bioaerosols in indoor air, *Ann. Agric. Environ. Med.* 6 (1999) 105–113.
- [59] B.F. Habbick, M.M. Pizzichini, B. Taylor, D. Rennie, A. Senthilselvan, M.R. Sears, Prevalence of asthma, rhinitis and eczema among children in 2 Canadian cities: the international study of asthma and allergies in childhood, *Can. Med. Assoc. J.* 160 (13) (1999) 1824–1828.
- [60] T.D. Hartwell, R.L. Perritt, E.D. Pellizzari, L.C. Michael, Results from the 1987 total exposure assessment methodology (TEAM) study in southern California, *Atmos. Environ.* 26A (1992) 1519–1527.
- [61] A.L. Heffernan, P. Sly, L. Hearn, N. Holling, J.F. Mueller, The bedroom: an exposure source for polybrominated flame retardants (PBDEs) in infants?, in: *Proceedings of Dioxin 2011*, 2011. Brussels, Belgium.
- [62] J. Heyes, R. Siebers, F.I. Biol, A. Parkes, N. Piersie, J. Crane,  $\beta$ -(1,3)-Glucan levels and its determinants in New Zealand bedrooms, *J. Asthma* 46 (2009) 64–66.
- [63] J.B. Hicks, E.T. Lu, R. De Guzman, M. Weingart, Fungal types and concentrations from settled dust in normal residences, *J. Occup. Environ. Hyg.* 2 (2005) 481–492.
- [64] K. Hillier, T. Schupp, I. Carney, An investigation into VOC emissions from polyurethane flexible foam mattresses, *Cell. Polym.* 22 (4) (2003) 237–259.
- [65] K. Hillier, D.A. King, C. Henneuse, Study of odours coming out of polyurethane flexible foam mattresses, *Cell. Polym.* 28 (2) (2009) 113–144.
- [66] D. Hospodsky, J. Qian, W.W. Nazaroff, N. Yamamoto, K. Bibby, H. Rismani-Yazdi, J. Peccia, Human occupancy as a source of indoor airborne bacteria, *PLoS One* 7 (2012) e34867.
- [67] M.G. Howat, Postulates for partitioning variance. I., in body movement among its origins, *J. Theor. Biol.* 37 (1972) 125–137.
- [68] N.-Y. Hsu, C.-Y. Chen, C.-C. Lee, H.-J. Su, Relationship between Indoor Phthalate Concentrations and Dampness or Visible Mold, in: *Proceedings of Healthy Buildings 2012*, 2012. Brisbane, Australia, paper ID: 7B.4.
- [69] H.-M. Hwang, E.-K. Park, T.M. Young, B.D. Hammock, Occurrence of endocrine-disrupting chemicals in indoor dust, *Sci. Total Environ.* 404 (2008) 26–35.
- [70] I. Iglowstein, O.G. Jenni, L. Molinari, R.H. Largo, Sleep duration from infancy to adolescence: reference values and generational trends, *Pediatrics* 111 (2003) 302–307.
- [71] C. Instanes, G. Hetland, S. Berntsen, M. Løvik, P. Nafstad, Allergens and endotoxin in settled dust from day-care centers and schools in Oslo, Norway, *Indoor Air* 15 (2005) 356–362.
- [72] P.A. Jensen, D. O'Brien, Industrial hygiene, in: K. Willeke, P.A. Baron (Eds.), *Aerosol Measurement: Principles, Techniques, and Applications*, Van Nostrand Reinhold, New York, NY, 1993, pp. 537–559.
- [73] S. Jovanovic, A. Felder-Kennel, T. Gabrio, B. Kouros, B. Link, V. Maisner, I. Piechotowski, K.H. Schick, M. Schimpf, U. Weidner, I. Zöllner, M. Schwenk, Indoor fungi levels in homes of children with and without allergy history, *Int. J. Hyg. Environ. Health* 207 (2004) 369–378.
- [74] A.S. Keller, N.P. Raju, T.F. Webster, H.M. Stapleton, Flame retardant applications in camping tents and potential exposure, *Environ. Sci. Technol. Lett.* 1 (2) (2014) 152–155.
- [75] S. Kemmlin, O. Hahn, O. Jann, Emissions of organophosphate and brominated flame retardants from selected consumer products and building materials, *Atmos. Environ.* 37 (2003) 5485–5493.
- [76] S. Kilburn, T.J. Lasserson, M.P. McKean, Pet allergen control measures for allergic asthma in children and adults, *Cochrane Database Syst. Rev.* 1 (2003) CD002989.
- [77] N.E. Klepeis, W.C. Nelson, W.R. Ott, J.P. Robinson, A.M. Tsang, P. Switzer, J.V. Behar, S.C. Hern, W.H. Engelmann, The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants, *J. Expo. Anal. Environ. Epidemiol.* 11 (2001) 231–252.
- [78] H.S.M. Kort, G. Schober, L.G.H. Koren, J. Scharringa, Mould-devouring mites differ in guanine excretion from dust-eating Acari, a possible error source in mite allergen exposure studies, *Clin. Exp. Allergy* 27 (1997) 921–925.
- [79] M. Korthals, M. Ege, S. Lick, E. von Mutius, J. Bauer, Occurrence of *Listeria* spp. in mattress dust of farm children in Bavaria, *Environ. Res.* 107 (2008) 299–394.
- [80] U. Krämer, C. Lemmen, E. Bartusel, E. Link, J. Ring, H. Behrendt, Current eczema in children is related to Der f 1 exposure but not to Der p 1 exposure, *Brit. J. Dermatol* 154 (2006) 99–105.
- [81] C.A. Krone, J.T.A. Ely, T. Klinger, R.J. Rando, Isocyanates in flexible polyurethane foams, *Bull. Environ. Contam. Toxicol.* 70 (2003) 328–335.
- [82] E.J.M. Krop, G. Doekes, M.J. Stone, R.C. Aalberse, J.S. van der Zee, Spreading of occupational allergens: laboratory animal allergens on hair-covering caps and in mattress dust of laboratory animal workers, *Occup. Environ. Med.* 64 (2007) 267–272.
- [83] I. Kulmala, A. Säämänen, S. Enbom, The effect of contaminant source location on worker exposure in the near-wake region, *Ann. Occup. Hyg.* 40 (1996) 511–523.
- [84] L. Lan, Z. Lian, X. Zhou, C. Sun, H. Huang, Y. Lin, J. Zhao, Pilot study on the application of bedside personalized ventilation to sleeping people, *Build. Environ.* 67 (2013) 160–166.
- [85] P.J. Landrigan, L.R. Goldman, Children's vulnerability to toxic chemicals: a challenge and opportunity to strengthen health and environmental policy, *Health Aff.* 30 (5) (2011) 842–850.
- [86] B.P. Lanphear, C.V. Vorhees, D.C. Bellinger, Protecting children from environmental toxins, *PLoS Med.* 2 (3) (2005) e61.
- [87] J. Laverge, A. Janssens, Analysis of the Influence of Ventilation Rate on Sleep Pattern, in: *Proceedings of Indoor Air 2011*, 2011. Austin, Texas, USA, paper ID 51.
- [88] J. Laverge, A. Novoselac, R.L. Corsi, A. Janssens, Experimental assessment of exposure to gaseous pollutants from mattresses and pillows while asleep, *Build. Environ.* 59 (2013) 203–210.
- [89] J. Laverge, M.P. Spilak, A. Novoselac, Experimental assessment of the inhalation zone of standing, sitting and sleeping persons, *Build. Environ.* 82 (2015a) 258–266.
- [90] J. Laverge, M. Delghust, A. Janssens, Carbon dioxide concentrations and humidity levels measured in Belgian standard and low energy dwellings with common ventilation strategies, *Inter. J. Vent.* 14 (2) (2015b) 165–180.
- [91] J.A. Lawson, J.A. Dosman, D.C. Rennie, J.R. Beach, S.C. Newman, T. Crowe, A. Senthilselvan, Endotoxin as a determinant of asthma and wheeze among rural dwelling children and adolescents: a case-control study, *BMC Pulm. Med.* 12 (2012) 56.
- [92] J.A. Leech, K. Wilby, E. McMullen, K. Laporte, The Canadian human activity pattern survey: report of methods and population surveyed, *Chron. Dis. Can.* 17 (1996) 118–123.
- [93] T.F. Leung, Y.S. Wong, I.H.S. Chan, E. Yung, H.Y. Sy, C.W.K. Lam, G.W.K. Wong, Domestic exposure to aeroallergens in Hong Kong families with asthmatic children, *Pediatr. Pulmonol.* 46 (2011) 632–639.
- [94] Y. Liang, Y. Xu, Improved method for measuring and characterizing phthalate emissions from building materials and its application to exposure assessment, *Environ. Sci. Technol.* 48 (8) (2014a) 4475–4484.
- [95] Y. Liang, Y. Xu, Emission of phthalates and phthalate alternatives from vinyl flooring and crib mattress covers: the influence of temperature, *Environ. Sci. Technol.* 48 (24) (2014b) 14228–14237.
- [96] D. Licina, Y. Tian, W.W. Nazaroff, Emission rates and the personal cloud effect associated with particle release from the perihuman environment, *Indoor Air* (2016), <http://dx.doi.org/10.1111/ina.12365>.
- [97] G. Lidén, J. Waher, Experimental investigation of the concept of a 'breathing zone' using a mannequin exposed to a point source of inertial/sedimenting particles emitted with momentum, *Ann. Occup. Hyg.* 54 (2010) 100–116.
- [98] C.-C. Lin, K.-P. Yu, P. Zhao, G.W.-M. Lee, Evaluation of impact factors on VOC emissions and concentrations from wooden flooring based on chamber tests, *Build. Environ.* 44 (2009) 525–533.
- [99] L. Lu, T. Tamura, T. Togawa, Detection of body movements during sleep by monitoring of bed temperature, *Physiol. Meas.* 20 (1999) 137–148.
- [100] T. Madsen, R. Gibson, Toxic Baby Furniture: the Latest Case for Making Products Safe from the Start, *Environmental California Research & Policy Center*, 2008.
- [101] D.T. Mage, W.R. Ott, Accounting for nonuniform mixing and human exposure in indoor environments, in: Bruce A. Tichenor (Ed.), *Characterizing Sources of Indoor Air Pollution and Related Sink Effects*, ASTM STP 1287, American Society for Testing and Materials, 1996, pp. 263–278.
- [102] A. Mahmic, E. Tovey, C.A. Molloy, L. Young, House dust mite allergen exposure in infancy, *Clin. Exp. Allergy* 28 (1998) 1487–1492.
- [103] S. McBride, A.R. Ferro, W.R. Ott, P. Switzer, L.M. Hildemann, Investigations of the proximity effects for pollutants in the indoor environment, *J. Expo. Anal. Environ. Epidemiol.* 9 (1999) 602–621.
- [104] J.F. Meadow, A.E. Altrichter, A.C. Bateman, J. Stenson, G.Z. Brown, J.L. Green, B.J.M. Bohannon, Humans differ in their personal microbial cloud, *PeerJ.* 3 (2015) e1258.
- [105] T. Meklin, T. Reponen, M. Toivola, V. Koponen, T. Husman, A. Hyvärinen, A. Nevalainen, Size distributions of airborne microbes in moisture-damaged and reference school buildings of two construction types, *Atmos. Environ.* 36 (2002) 6031–6039.
- [106] A.K. Melikov, Z.D. Bolashikov, E. Georgiev, Novel ventilation strategy for reducing the risk of airborne cross infection in hospital rooms, in: *Proceedings of Indoor Air 2011*, Austin, Texas, USA, Paper ID 1037, 2011.
- [107] S. Mihrshahi, G. Marks, C. Vanlaar, E. Tovey, J. Peat, Predictors of high house dust mite allergen concentrations in residential homes in Sydney, *Allergy* 57 (2002) 137–142.
- [108] G. Morrison, H. Li, S. Mishra, M. Buechlein, Airborne phthalate partitioning to cotton clothing, *Atmos. Environ.* 115 (2015) 149–152.
- [109] G. Morrison, C.J. Weschler, G. Bekö, Dermal uptake directly from air under transient conditions: advances in modeling and comparisons with experimental results for human subjects, *Indoor Air* 26 (2016) 913–924.
- [110] H.-S. Nam, R. Siebers, S.-H. Lee, J.-S. Park, Y.-B. Kim, Y.-J. Choi, S.-H. Lee, J. Crane, House dust mite allergens in domestic homes in Cheonan, Korea, *Korean J. Parasitol* 46 (2008) 187–189.
- [111] National Sleep Foundation (NSF), 2010 Sleep in America Poll: Summary of Findings, 2010. Accessed online: <https://sleepfoundation.org/sites/default/>

- files/nsaw/NSF%20Sleep%20in%20America%20Poll%20-%20Summary%20of%20Findings%20.pdf.
- [112] M. Nnadili, N.E. Crain, R.L. Corsi, Effects of Relative Humidity on Chemical Off-gassing in Residences, in: Proceedings of Indoor Air 2011, 2011. Austin, Texas, USA paper ID 1147.
- [113] I. Odeh, T. Hussein, Activity pattern of urban adult students in an eastern mediterranean society, *Int. J. Environ. Res. Public Health* 13 (10) (2016) 960–965.
- [114] OECD, How Do People in the Asia/pacific Region Spend Their Time? Society at a Glance, Asia/Pacific 2011, 2011.
- [115] OEKO-TEX, Standard 100 by the International Association for Research and Testing in the Field of Textile and Leather Ecology. Edition 02.2017, 2017.
- [116] T. O'Meara, E. Tovey, Monitoring personal allergen exposure, *Clin. Rev. Allergy Immunol.* 18 (2000) 341–395.
- [117] H. Oppermann, C. Doering, A. Sobottka, U. Krämer, B. Thriene, Comparison of East and West German households with regard to exposure to house dust mites and mould fungi, *Gesundh. Wes* 63 (2001) 85–89 (in German).
- [118] C. Pedroletti, E. Millinger, B. Dahlen, P. Söderman, O. Zetterström, Clinical effects of purified air administered to the breathing zone in allergic asthma: a double-blind randomized cross-over trial, *Resp. Med.* 103 (9) (2009) 1313–1319.
- [119] F.-A. Pitten, M. Scholler, U. Krüger, I. Effendy, A. Kramer, Filamentous fungi and yeasts on mattresses covered with different encasings, *Eur. J. Dermatol* 11 (2001) 534–537.
- [120] T. Platts-Mills, P. Heymann, J. Longbottom, S. Wilkins, Airborne allergens associated with asthma: particle sizes carrying dust mite and rat allergens measured with a cascade impactor, *J. Allergy Clin. Immunol.* 77 (1986) 850–857.
- [121] J. Qian, J. Peccia, A.R. Ferro, Walking-induced particle resuspension in indoor environments, *Atmos. Environ.* 89 (2014) 464–481.
- [122] N. Rabinovitch, A.H. Liu, L. Zhang, C.E. Rodes, K. Foarde, S.J. Dutton, J.R. Murphy, E.W. Gelfand, Importance of the personal endotoxin cloud in school-age children with asthma, *J. Allergy Clin. Immunol.* 116 (2005) 1053–1057.
- [123] S. Raja, Y. Xu, A.R. Ferro, P.A. Jaques, P.K. Hopke, Resuspension of indoor aeroallergens and relationship to lung inflammation in asthmatic children, *Environ. Inter* 36 (2010) 8–14.
- [124] Rakennustieto, M1 Emission Classification of Building Materials: Protocol for Chemical and Sensory Testing of Building Materials. M1 Emission Class for Building Materials, 2017. Version 5.7.2017.
- [125] D.C. Rennie, J.A. Lawson, S.P. Kirychuk, C. Paterson, P.J. Willson, A. Senthilselvan, D.W. Cockcroft, Assessment of endotoxin levels in the home and current asthma and wheeze in school-age children, *Indoor Air* 18 (2008) 447–453.
- [126] T. Reponen, S.A. Grinshpun, K.L. Conwell, J. Wiest, M. Anderson, Aerodynamic versus physical size of spores: measurement and implication for respiratory deposition, *Grana* 40 (2001) 119–125.
- [127] T. Reponen, S.-C. Seo, F. Grimley, T. Lee, C. Crawford, S.A. Grinshpun, Fungal fragments in moldy homes: a field study in homes in New Orleans and Southern Ohio, *Atmos. Environ.* 41 (2007) 8140–8149.
- [128] D. Rim, A. Novoselac, Transport of particulate and gaseous pollutants in the vicinity of a human body, *Build. Environ.* 44 (2009) 1840–1849.
- [129] D. Rim, A. Novoselac, Occupational exposure to hazardous airborne pollutants: effects of source characteristics, *J. Occup. Environ. Hyg.* 7 (2010) 683–692.
- [130] C.E. Rodes, R.M. Kamens, R.W. Wiener, The significance and characteristics of the personal activity cloud on exposure assessment measurements for indoor air, *Indoor Air* 2 (1991) 123–145.
- [131] C.E. Rodes, P.A. Lawless, G.F. Evans, L.S. Sheldon, R.W. Williams, A.F. Vette, J.P. Creason, D. Walsh, The relationships between personal PM exposures for elderly populations and indoor and outdoor concentrations for three retirement center scenarios, *J. Expo. Anal. Environ. Epidemiol.* 11 (2001) 103–115.
- [132] C.E. Rodes, Indoor aerosol exposure assessment, in: P. Kulkarni, P.A. Baron, K. Willeke (Eds.), *Aerosol Measurement: Principles, Techniques, and Applications*, John Wiley & Son, Inc., New York, NY, 2011, pp. 615–634.
- [133] R.A. Rudel, L.J. Perovich, Endocrine disrupting compounds in indoor and outdoor air, *Atmos. Environ.* 43 (2009) 170–181.
- [134] M. Sakaguchi, S. Inouye, H. Yasueda, T. Shida, Concentration of airborne mite allergens (Der I and Der II) during sleep, *Allergy* 47 (1992) 55–57.
- [135] T. Salthammer, F. Fuhrmann, E. Uhde, Flame retardants in the indoor environment – Part II: release of VOCs (triethylphosphate and halogenated degradation products) from polyurethane, *Indoor Air* 13 (2003) 49–52.
- [136] U. Schauer, K.C. Bergmann, M. Gerstlauer, S. Lehmann, M. Gappa, A. Brenneken, C. Schulz, P. Ahrens, J. Schreiber, M. Wittmann, E. Hamelmann, Improved asthma control in patients with severe, persistent allergic asthma after 12 months of nightly temperature-controlled laminar airflow: an observational study with retrospective comparisons, *Eur. Clin. Respir. J.* 2 (2015), <http://dx.doi.org/10.3402/ecrj.v2.28531>.
- [137] C. Schweizer, R.D. Edwards, L. Bayer Oglesb, W.J. Gauderman, V. Ilacqua, M.J. Jantunen, H.K. Lai, M. Nieuwenhuijsen, N. Künzli, Indoor time-microenvironment activity patterns in seven regions of Europe, *J. Expo. Sci. Environ. Epidemiol.* 17 (2007) 170–181.
- [138] K.E. Sidenius, T.E. Hallas, J. Stenderup, L.K. Poulsen, H. Mosbech, Decay of house-dust mite allergen Der f 1 at indoor climatic conditions, *Ann. Allergy Asthma Immunol.* 89 (1) (2002) 34–37.
- [139] M. Soleimani, J. Rafinejad, House dust mite contamination in hotels and inns in Bandar Abbas, South of Iran, Iran. *J. Environ. Health Sci. Eng.* 5 (2008) 207–210.
- [140] F. Spertini, M. Berney, F. Foradini, C.-A. Roulet, Major mite allergen Der f 1 concentration is reduced in buildings with improved energy performance, *Allergy* 65 (2009) 623–629.
- [141] M.P. Spilak, B.E. Boor, A. Novoselac, R.L. Corsi, Impact of bedding arrangements, pillows, and blankets on particle resuspension in the sleep micro-environment, *Build. Environ.* 81 (2014a) 60–68.
- [142] M.P. Spilak, G.D. Karottki, B. Kolarik, M. Frederiksen, S. Loft, L. Gunnarsen, Evaluation of building characteristics in 27 dwellings in Denmark and the effect of using particle filtration units on PM 2.5 concentrations, *Build. Environ.* 73 (2014b) 55–63.
- [143] M.P. Spilak, T. Sigsgaard, H. Takai, G. Zhang, A comparison between temperature-controlled laminar airflow device and a room air-cleaner in reducing exposure to particles while asleep, *PLoS One* 11 (11) (2016) e0166882.
- [144] H.M. Stapleton, S. Klosterhaus, S. Eagle, J. Fuh, J.D. Meeker, A. Blum, T.F. Webster, Detection of organophosphate flame retardants in furniture foam and U.S. house dust, *Environ. Sci. Technol.* 43 (2009) 7490–7495.
- [145] H.M. Stapleton, S. Klosterhaus, A. Keller, P. Lee Ferguson, S. van Bergen, E. Cooper, T.F. Webster, A. Blum, Identification of flame retardants in polyurethane foam collected from baby products, *Environ. Sci. Technol.* 45 (2011) 5323–5331.
- [146] M.M. Stein, C.L. Hrusch, J. Gozdz, et al., Innate immunity and asthma risk in Amish and Hutterite farm children, *N. Engl. J. Med.* 375 (2016) 411–421.
- [147] B. Stephens, J.A. Siegel, A. Novoselac, Operational characteristics of residential and light-commercial air-conditioning systems in a hot and humid climate zone, *Build. Environ.* 46 (2011) 1972–1983.
- [148] P. Strøm-Tejsten, D. Zukowska, P. Wargocki, D.P. Wyon, The effects of bedroom air quality on sleep and next-day performance, *Indoor Air* 26 (2016) 679–686.
- [149] A.H. Stubner, K.H. Dillon, C.L. Kohler, Home remediation for respiratory health: a feasibility study, *Fam. Community Health* 22 (2000) 1–15.
- [150] H.-J. Su, P.-C. Wu, H.-L. Chen, F.-C. Lee, L.-L. Lin, Exposure assessment of indoor allergens, endotoxin, and airborne fungi for homes in southern Taiwan, *Environ. Res. Sec. A* 85 (2001) 135–144.
- [151] M. Täubel, H. Rintala, M. Pitkäranta, L. Paulin, S. Laitinen, J. Pekkanen, A. Hyvärinen, A. Nevalainen, The occupant as a source of house dust bacteria, *J. Allergy Clin. Immunol.* 124 (2009) 834–840.
- [152] Y. Tian, K. Sul, J. Qian, S. Mondal, A.R. Ferro, A comparative study of walking-induced dust resuspension using a consistent test mechanism, *Indoor Air* 24 (2014) 592–603.
- [153] C. Tischer, U. Gehring, C.M. Chen, M. Kerkhof, G. Koppelman, S. Sausenthaler, O. Herbarth, B. Schaaf, I. Lehmann, U. Krämer, D. Berdel, A. von Berg, C.P. Bauer, S. Koletzko, H.E. Wichmann, B. Brunekreef, J. Heinrich, Respiratory health in children, and indoor exposure to (1,3)-β-D-glucan, EPS mould components and endotoxin, *Eur. Respir. J.* 37 (2011) 1050–1059.
- [154] C. Tischer, J.-P. Zock, M. Valkonen, G. Doekes, S. Guerra, D. Heederik, D. Jarvis, D. Norbäck, M. Olivieri, J. Sunyer, C. Svanes, M. Täubel, E. Thiering, G. Verlato, A. Hyvärinen, J. Heinrich, Predictors of microbial agents in dust and respiratory health in the Ecrhs, *BMC Pulm. Med.* 15 (2015) 48.
- [155] L.-M.L. Toms, L. Hearn, K. Kennedy, F. Harden, M. Bartkow, C. Temme, J.F. Mueller, Concentrations of polybrominated diphenyl ethers (PBDEs) in matched samples of human milk, dust, and indoor air, *Environ. Int.* 35 (2009) 864–869.
- [156] E.R. Tovey, D. Liu-Brennan, F.L. Garden, B.G. Oliver, M.S. Perzanowski, G.B. Marks, Time-based measurement of personal mite allergen bioaerosol exposure over 24 hour periods, *PLoS ONE* 11 (5) (2016) e0153414.
- [157] UL Environment Standard, GREENGUARD Certification Program Method for Measuring and Evaluating Chemical Emissions from Building Materials, Finishes and Furnishings. Standard 2821, Edition 1, 2013.
- [158] U.S. Consumer Product Safety Commission (CPSC), Consumer Product Safety Improvement Act (CPSIA) of 2008, U.S. CPSC, Bethesda, Maryland, 2008.
- [159] U.S. Environmental Protection Agency (EPA), Exposure Factors Handbook, Chapter 6, Inhalation Rates, U.S. EPA, Washington, D.C., 2009.
- [160] M. Valkonen, I.M. Wouters, M. Täubel, H. Rintala, V. Lenters, R. Vasara, J. Genuneit, C. Braun-Fahrländer, R. Piarroux, E. von Mutius, D. Heederik, A. Hyvärinen, Bacterial exposures and associations with atopy and asthma in children, *PLoS ONE* 10 (6) (2015) e0131594.
- [161] L. van den Bemt, M.P. de Vries, L. van Knapen, M. Jansen, M. Goossens, J.W.M. Muris, C.P. van Schayck, Influence of mattress characteristics on house dust mite allergen concentration, *Clin. Exp. Allergy* 36 (2006) 233–237.
- [162] E. Vangronsveld, S. Berckmans, M. Spence, Toluene diisocyanate emission to air and migration to a surface from a flexible polyurethane foam, *Ann. Occup. Hyg.* 57 (5) (2013) 650–661.
- [163] N. Visitsunthorn, V. Chirdjirapong, V. Pootong, O. Jirapongsananuruk, P. Pacharn, S. Weeravejsukit, V. Mahakittikun, P. Vichyanond, The accumulation of dust mite allergens on mattresses made of different kinds of materials, *Asian pac. J. Allergy Immunol.* 28 (2010) 155–161.
- [164] K. Vogel, N. Blümer, M. Korthals, J. Mittelstädt, H. Garn, M. Ege, E. von Mutius, S. Gatermann, A. Bufe, T. Goldmann, K. Schwaiger, H. Renz, S. Brandau, J. Bauer, H. Heine, O. Holst, Animal shed *Bacillus licheniformis* spores possess allergy-protective as well as inflammatory properties, *J. Allergy Clin.*

- Immunol. 122 (2008) 307–312.
- [165] L.A. Wallace, Indoor particles: a review, *J. Air Waste Manag. Assoc.* 46 (1996) 98–126.
- [166] L. Wang, Q. Chen, Evaluation of some assumptions used in multizone airflow network models, *Build. Environ.* 43 (2007) 1671–1677.
- [167] C.J. Weschler, W.W. Nazaroff, Semivolatile organic compounds in indoor environment, *Atmos. Environ.* 42 (2008) 9018–9040.
- [168] C.J. Weschler, S. Langer, A. Fischer, G. Bekö, J. Toftum, G. Clausen, Squalene and cholesterol in dust from Danish homes and daycare centers, *Environ. Sci. Technol.* 45 (2011) 3872–3879.
- [169] C.J. Weschler, W.W. Nazaroff, SVOC exposure indoors: fresh look at dermal pathways, *Indoor Air* 22 (2012) 356–377.
- [170] C.J. Weschler, G. Bekö, H.M. Koch, T. Salthammer, T. Schripp, J. Toftum, G. Clausen, Transdermal uptake of diethyl phthalate and di(n-butyl) phthalate directly from air: experimental verification, *Environ. Health Persp* 123 (10) (2015) 928–934.
- [171] A.A. Woodcock, N. Steel, C.B. Moore, S.J. Howard, A. Custovic, D.W. Denning, Fungal contamination of bedding, *Allergy* 61 (2006) 140–142.
- [172] F.F.-S. Wu, R. Siebers, F.I. Biol, C.-F. Chang, S.-W. Hsieh, M.-W. Wu, C.-Y. Chen, N. Pierse, J. Crane, Indoor allergens and microbial bio-contaminants in homes of asthmatic children in central Taiwan, *J. Asthma* 46 (2009) 745–749.
- [173] F.F.-S. Wu, M.-W. Wu, N. Pierse, J. Crane, R. Siebers, Daily vacuuming of mattresses significantly reduces house dust mite allergens, bacterial endotoxin, and fungal  $\beta$ -glucan, *J. Asthma* 49 (2012) 139–143.
- [174] Y. Xu, J.C. Little, Predicting emissions of SVOCs from polymeric materials and their interaction with airborne particles, *Environ. Sci. Technol.* 40 (2) (2006) 456–461.
- [175] Y. Xu, E.A. Cohen Hubal, J.C. Little, Predicting residential exposure to phthalate plasticizer emitted from vinyl flooring: sensitivity, uncertainty, and implications for biomonitoring, *Environ. Health Persp* 118 (2010) 253–258.
- [176] Y. Xu, Z. Liu, J. Park, P.A. Clausen, J.L. Benning, J.C. Little, Measuring and predicting the emission rate of phthalate plasticizer from vinyl flooring in a specially-designed chamber, *Environ. Sci. Technol.* 46 (22) (2012) 12534–12541.
- [177] W. Yang, K. Lee, C. Yoon, S. Yu, K. Park, W. Cho, Determinants of residential indoor and transportation activity times in Korea, *J. Expo. Anal. Environ. Epidemiol.* 21 (2011) 310–316.
- [178] X. Yang, Q. Chen, J.S. Zhang, Impact of early stage incomplete mixing on estimating VOC emissions in small test chambers, *Indoor Air* 8 (1998) 180–189.
- [179] V.G. Zartarian, W.R. Ott, N. Duan, A quantitative definition of exposure and related concepts, *J. Expo. Anal. Env. Epid* 7 (1997) 411–437.
- [180] A.R. Ferro, R.J. Kopperud, L.M. Hildemann, Elevated personal exposure to particulate matter from human activities in a residence, *J. Expo. Anal. Env. Epid.* 14 (2004) S34–S40.
- [181] C.J. Weschler, W.W. Nazaroff, Dermal uptake of organic vapors commonly found in indoor air, *Environ. Sci. Technol.* 48 (2) (2013) 1230–1237.
- [182] E.A. Cohen Hubal, M.G. Nishioka, W.A. Ivancic, M. Morara, P.P. Eggeghy, Comparing surface residue transfer efficiencies to hands using polar and nonpolar fluorescent tracers, *Environ. Sci. Technol.* 42 (3) (2008) 934–939.
- [183] H.F. Frasch, G.S. Dotson, A.L. Bunge, C.P. Chen, J.W. Cherrie, G.B. Kasting, J.C. Kissel, J. Sahlmeier, S. Semple, S. Wilkinson, Analysis of finite dose dermal absorption data: Implications for dermal exposure assessment, *J. Expo. Anal. Environ. Epid.* 24 (1) (2014) 65–73.
- [184] U.S. Census Bureau, United States Census 2010, 2010. <https://www.census.gov/2010census/>.
- [185] J.M. Hugo, M.W. Spence, T.D. Lickly, The determination of the ability of polyurethane foam to release toluene diisocyanate into air, *Appl. Occup. Environ. Hyg.* 15 (6) (2000) 512–519.
- [186] S.T. Aaronson, S. Rashed, M.P. Biber, J.A. Hobson, Brain state and body position, *Arch. Gen. Psychiat.* 39 (1982) 330–335.
- [187] K. Azumi, S. Shirikawa, S. Takahashi, A proposal of a new classification for body movements during sleep: variations of body movements through a week, *Sleep Res.* 6 (1977) 44.
- [188] F. Giganti, G. Ficca, S. Gori, P. Salzarulo, Body movements during night sleep and their relationship with sleep stages are further modified in very old subjects, *Brain Res. Bull.* 75 (2008) 66–69.
- [189] M.H. Silber, S. Ancoli-Israel, M.H. Bonnet, S. Chokroverty, M.M. Grigg-Damberger, M. Hirshkowitz, S. Kapen, S.A. Keenan, M.H. Kryger, T. Penzel, M.R. Pressman, C. Iber, The visual scoring of sleep in adults, *J. Clin. Sleep Med.* 3 (2007) 121–131.
- [190] M. Shimohira, T. Shiiki, J. Sugimoto, Y. Ohsawa, M. Fukumizu, T. Hasegawa, Y. Iwakawa, Y. Nomura, M. Segawa, Video analysis of gross body movements during sleep, *Psychiatry Clin. Neurosci.* 52 (1998) 176–177.
- [191] J. Wilde-Frenz, H. Schulz, Rate and distribution of body movements during sleep in humans, *Percept. Mot. Skills* 56 (1983) 275–283.
- [192] C. Janson, W. De Backer, T. Gislason, P. Plaschke, E. Björnsson, J. Hetta, H. Kristbjarnarson, P. Vermeire, G. Boman, Increased prevalence of sleep disturbances and daytime sleepiness in subjects with bronchial asthma: a population study of young adults in three European countries, *Eur. Respir. J.* 9 (1996) 2132–2138.
- [193] G. Stores, A.J. Ellis, L. Wiggins, C. Crawford, A. Thomson, Sleep and psychological disturbance in nocturnal asthma, *Arch. Dis. Child.* 78 (1998) 413–419.
- [194] A. Zanobetti, S. Redline, J. Schwartz, D. Rosen, S. Patel, G.T. O'Connor, M. Lebowitz, B.A. Coull, D.R. Gold, Associations of PM10 with sleep and sleep-disordered breathing in adults from seven U.S. urban areas, *Am. J. Respir. Crit. Care Med.* 182 (2010) 819–825.
- [195] H. Homma, M. Yakiyama, Examination of free convection around occupant's body caused by its metabolic heat, *ASHRAE Trans.* 94 (1988) 104–124.
- [196] D. Zukowska, A. Melikov, Z. Popiolek, Impact of personal factors and furniture arrangement on the thermal plume above a sitting occupant, *Build. Environ.* 49 (2012) 104–116.
- [197] F. Haghghat, L.D. De Bellis, Material emission rates: literature review, and the impact of indoor air temperature and relative humidity, *Build. Environ.* 33 (1998) 261–277.
- [198] P.A. Clausen, Z. Liu, V. Kofoed-Sorensen, J. Little, P. Wolkoff, Influence of temperature on the emission of Di-(2-ethylhexyl)phthalate (DEHP) from PVC flooring in the emission cell FLEC, *Environ. Sci. Technol.* 46 (2) (2012) 909–915.
- [199] N. Canha, J. Lage, S. Candeias, C. Alves, S.M. Almeida, Indoor air quality during sleep under different ventilation patterns, *Atmos. Pollut. Res.* 25 (May 2017) in press, <https://doi.org/10.1016/j.apr.2017.05.004>.
- [200] T. Hirsch, M. Hering, K. Bürkner, D. Hirsch, W. Leupold, M.-L. Kerkmann, E. Kuhlisch, L. Jatzwauk, House-dust-mite allergen concentrations (Der f 1) and mold spores in apartment bedrooms before and after installation of insulated windows and central heating systems, *Allergy* 55 (2000) 79–83.
- [201] National Resources Defense Council (NRDC). Letter to U.S. CPSC. Re: Notice of Availability of Draft Guidance Regarding Which Children's Products are Subject to the Requirements of CPSIA Section 108; Request for Comments and Information. [http://www.nrdc.org/health/files/hea\\_09032501a.pdf](http://www.nrdc.org/health/files/hea_09032501a.pdf).